

BRCC36 Rabbit mAb

Catalog # AP75167

Product Information

Application	WB, FC
Primary Accession	P46736
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	36072

Additional Information

Gene ID	79184
Other Names	BRCC3
Dilution	WB~~1:1000-1:5000 FC~~1:100-1:500
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	BRCC3
Synonyms	BRCC36, C6.1A, CXorf53
Function	Metalloprotease that specifically cleaves 'Lys-63'-linked polyubiquitin chains (PubMed: 19214193 , PubMed: 20656690 , PubMed: 24075985 , PubMed: 26344097). Does not have activity toward 'Lys- 48'-linked polyubiquitin chains (PubMed: 19214193 , PubMed: 20656690 , PubMed: 24075985 , PubMed: 26344097). Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs) (PubMed: 14636569 , PubMed: 16707425 , PubMed: 17525341 , PubMed: 19202061 , PubMed: 19261746 , PubMed: 19261748 , PubMed: 19261749). In the BRCA1-A complex, it specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX, antagonizing the RNF8-dependent ubiquitination at double-strand breaks (DSBs)

(PubMed:[20656690](#)). Catalytic subunit of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates (PubMed:[20656690](#), PubMed:[24075985](#), PubMed:[26195665](#), PubMed:[26344097](#)). Mediates the specific 'Lys-63'-specific deubiquitination associated with the COP9 signalosome complex (CSN), via the interaction of the BRISC complex with the CSN complex (PubMed:[19214193](#)). The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1 (PubMed:[26195665](#)). Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression (PubMed:[24075985](#), PubMed:[26344097](#)). Acts as a regulator of the NLRP3 inflammasome by mediating deubiquitination of NLRP3, leading to NLRP3 inflammasome assembly (By similarity). Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination (PubMed:[24075985](#)). Deubiquitinates HDAC1 and PWWP2B leading to their stabilization (By similarity).

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole Note=Localizes at sites of DNA damage at double-strand breaks (DSBs) (PubMed:[20656690](#), PubMed:[26344097](#)). Interaction with ABRAXAS2 retains BRCC3 in the cytoplasm (PubMed:[20656690](#)).

Tissue Location



Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Aberrantly expressed in the vast majority of breast tumors.

Background

Metalloprotease that specifically cleaves 'Lys-63'-linked polyubiquitin chains (PubMed:[19214193](#), PubMed:[20656690](#), PubMed:[24075985](#), PubMed:[26344097](#)). Does not have activity toward 'Lys-48'-linked polyubiquitin chains. Component of the BRCA1-A complex, a complex that specifically recognizes 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). In the BRCA1-A complex, it specifically removes 'Lys-63'-linked ubiquitin on histones H2A and H2AX, antagonizing the RNF8-dependent ubiquitination at double-strand breaks (DSBs) (PubMed:[20656690](#)). Catalytic subunit of the BRISC complex, a multiprotein complex that specifically cleaves 'Lys-63'-linked ubiquitin in various substrates (PubMed:[20656690](#), PubMed:[24075985](#), PubMed:[26344097](#), PubMed:[26195665](#)). Mediates the specific 'Lys-63'-specific deubiquitination associated with the COP9 signalosome complex (CSN), via the interaction of the BRISC complex with the CSN complex (PubMed:[19214193](#)). The BRISC complex is required for normal mitotic spindle assembly and microtubule attachment to kinetochores via its role in deubiquitinating NUMA1 (PubMed:[26195665](#)). Plays a role in interferon signaling via its role in the deubiquitination of the interferon receptor IFNAR1; deubiquitination increases IFNAR1 activity by enhancing its stability and cell surface expression (PubMed:[24075985](#), PubMed:[26344097](#)). Down-regulates the response to bacterial lipopolysaccharide (LPS) via its role in IFNAR1 deubiquitination (PubMed:[24075985](#)).

Images

Western blot analysis of BRCC36 in rat Brain, C6, 3T3, Hela lysates using BRCC36 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.