

CARD9 Rabbit mAb

Catalog # AP75188

Product Information

Application	WB, IP
Primary Accession	Q9H257
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	62241

Additional Information

Gene ID	64170
Other Names	CARD9
Dilution	WB~~1:1000-1:5000 IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	CARD9 {ECO:0000303 PubMed:11053425, ECO:0000312 HGNC:HGNC:16391}
Function	Adapter protein that plays a key role in innate immune response against fungi by forming signaling complexes downstream of C- type lectin receptors (PubMed: 26961233 , PubMed: 33558980). CARD9- mediated signals are essential for antifungal immunity against a subset of fungi from the phylum Ascomycota (PubMed: 24231284 , PubMed: 25057046 , PubMed: 25702837 , PubMed: 26521038 , PubMed: 26679537 , PubMed: 26961233 , PubMed: 27777981 , PubMed: 29080677 , PubMed: 33558980). Transduces signals in myeloid cells downstream of C-type lectin receptors CLEC7A (dectin-1), CLEC6A (dectin-2) and CLEC4E (Mincle), which detect pathogen- associated molecular pattern metabolites (PAMPs), such as fungal carbohydrates, and trigger CARD9 activation (By similarity). Upon activation, CARD9 homooligomerizes to form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10 and subsequent recruitment of MALT1: this leads to activation of NF-kappa-B and

MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed:[11053425](#), PubMed:[26488816](#), PubMed:[26961233](#), PubMed:[31296852](#), PubMed:[33558980](#)). CARD9 signaling in antigen-presenting cells links innate sensing of fungi to the activation of adaptive immunity and provides a cytokine milieu that induces the development and subsequent of interleukin 17-producing T helper (Th17) cells (PubMed:[24231284](#)). Also involved in activation of myeloid cells via classical ITAM- associated receptors and TLR: required for TLR-mediated activation of MAPK, while it is not required for TLR-induced activation of NF-kappa-B (By similarity). CARD9 can also be engaged independently of BCL10: forms a complex with RASGRF1 downstream of C-type lectin receptors, which recruits and activates HRAS, leading to ERK activation and the production of cytokines (By similarity). Acts as an important regulator of the intestinal commensal fungi (mycobiota) component of the gut microbiota (PubMed:[33548172](#)). Plays an essential role in antifungal immunity against dissemination of gut fungi: acts by promoting induction of antifungal IgG antibodies response in CX3CR1(+) macrophages to confer protection against disseminated *C.albicans* or *C.auris* infection (PubMed:[33548172](#)). Also mediates immunity against other pathogens, such as certain bacteria, viruses and parasites; CARD9 signaling is however redundant with other innate immune responses (By similarity). In response to *L.monocytogenes* infection, required for the production of inflammatory cytokines activated by intracellular peptidoglycan: acts by connecting NOD2 recognition of peptidoglycan to downstream activation of MAP kinases (MAPK) without activating NF- kappa-B (By similarity).

Cellular Location

Cytoplasm


Tissue Location

Expression is restricted to several populations of phagocytes, such as macrophages, monocytes, and dendritic cells (PubMed:[33548172](#)). Highly expressed in spleen (PubMed:[11053425](#)). Also detected in liver, placenta, lung, peripheral blood leukocytes and in brain (PubMed:[11053425](#)).


Background

Caspase recruitment domain-containing protein 9

Images



Western blot analysis of CARD9 in K562, THP-1 lysates using CARD9 antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.