

Chromogranin A Rabbit mAb

Catalog # AP75263

Product Information

Application WB, IHC-P, IHC-F, ICC

Primary Accession P10645
Reactivity Rat
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 50688

Additional Information

Gene ID 1113

Other Names CHGA

Dilution WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A ICC~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name CHGA

Function [Pancreastatin]: Strongly inhibits glucose induced insulin release from the

pancreas. [Serpinin]: Regulates granule biogenesis in endocrine cells by up-regulating the transcription of protease nexin 1 (SERPINE2) via a cAMP-PKA-SP1 pathway. This leads to inhibition of granule protein

degradation in the Golgi complex which in turn promotes granule formation.


Cellular Location [Serpinin]: Secreted {ECO:0000250 | UniProtKB:P26339}. Cytoplasmic vesicle,

secretory vesicle {ECO:0000250|UniProtKB:P26339}. Note=Pyroglutaminated serpinin localizes to secretory vesicle. {ECO:0000250|UniProtKB:P26339}

Tissue Location Detected in cerebrospinal fluid (at protein level) (PubMed:25326458).

Detected in urine (at protein level) (PubMed:37453717).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.