

Cyclin A2 Rabbit mAb

Catalog # AP75312

Product Information

Application WB, IHC-P, IHC-F, IP, ICC

Primary Accession P20248
Reactivity Human
Rabbit

Clonality Monoclonal Antibody

Calculated MW 48551

Additional Information

Gene ID 890

Other Names CCNA2

Dilution WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~N/A ICC~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name CCNA2 (HGNC:1578)

Function Cyclin which controls both the G1/S and the G2/M transition phases of the

cell cycle. Functions through the formation of specific serine/threonine protein kinase holoenzyme complexes with the cyclin- dependent protein kinases CDK1 or CDK2. The cyclin subunit confers the substrate specificity of these complexes and differentially interacts with and activates CDK1 and

CDK2 throughout the cell cycle.


Cellular Location Nucleus. Cytoplasm. Note=Exclusively nuclear during interphase


(PubMed:1312467). Detected in the nucleus and the cytoplasm at prophase


(PubMed:1312467). Cytoplasmic when associated with SCAPER

(PubMed:17698606).

Images







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.