

# Cytochrome C Rabbit mAb

Catalog # AP75324

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">P99999</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity Purified
<b>Calculated MW</b>	11749

## Additional Information

---

<b>Gene ID</b>	54205
<b>Other Names</b>	CYCS
<b>Dilution</b>	WB~~1:500-1:1000 IHC-P~~N/A
<b>Format</b>	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	CYCS
<b>Synonyms</b>	CYC
<b>Function</b>	Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain.
<b>Cellular Location</b>	Mitochondrion intermembrane space. Note=Loosely associated with the inner membrane


## Background

---



CYCS Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron

from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain.


## Images



Immunocytochemistry analysis of Cytochrome C (green) in K562 using Cytochrome C antibody, and DAPI (blue).



Western blot analysis of Cytochrome C in K562, rat Brain, C6, 3T3, HeLa lysates using Cytochrome C antibody.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.