

DDAH1 Rabbit mAb

Catalog # AP75346

Product Information

Application	WB, IHC-P, FC, IP
Primary Accession	O94760
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	31122

Additional Information

Gene ID	23576
Other Names	DDAH1
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A FC~~1:20-1:50 IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	DDAH1 (HGNC:2715)
Synonyms	DDAH
Function	Hydrolyzes N(G),N(G)-dimethyl-L-arginine (ADMA) and N(G)-monomethyl-L-arginine (MMA) which act as inhibitors of NOS. Has therefore a role in the regulation of nitric oxide generation.
Tissue Location	Detected in brain, liver, kidney and pancreas, and at low levels in skeletal muscle.

Background




Dimethylarginine dimethylaminohydrolase 1 (DDAH1) is an enzyme that can degrade asymmetric dimethylarginine (ADMA), an endogenous nitric oxide synthase (NOS)inhibitor (PMID:26996393). About 80%


of endogenous ADMA is metabolized by DDAH, principally the DDAH1 isoform (PMID:22460174). It was reported that DDAH1 is highly expressed in vascular endothelial cells in hearts (PMID:19917889). The observed molecular weight of DDAH1 is 35-40 kDa in the literature.

Images



Western blot analysis of DDAH1 in HeLa lysates using DDAH1 antibody.





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.