

# DYRK1A Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP7555c

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q13627</a>
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB12637
Calculated MW	85584
Antigen Region	389-418

## Additional Information

---

Gene ID	1859
Other Names	Dual specificity tyrosine-phosphorylation-regulated kinase 1A, Dual specificity YAK1-related kinase, HP86, Protein kinase minibrain homolog, MNBH, hMNB, DYRK1A, DYRK, MNB, MNBH
Target/Specificity	This DYRK1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 389-418 amino acids from the Central region of human DYRK1A.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DYRK1A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	DYRK1A {ECO:0000303 PubMed:25620562, ECO:0000312 HGNC:HGNC:3091}
Function	Dual-specificity kinase which possesses both serine/threonine and tyrosine kinase activities (PubMed: <a href="#">20981014</a> , PubMed: <a href="#">21127067</a> , PubMed: <a href="#">23665168</a> ,

PubMed:[30773093](#), PubMed:[8769099](#)). Exhibits a substrate preference for proline at position P+1 and arginine at position P-3 (PubMed:[23665168](#)). Plays an important role in double-strand breaks (DSBs) repair following DNA damage (PubMed:[31024071](#)). Mechanistically, phosphorylates RNF169 and increases its ability to block accumulation of TP53BP1 at the DSB sites thereby promoting homologous recombination repair (HRR) (PubMed:[30773093](#)). Also acts as a positive regulator of transcription by acting as a CTD kinase that mediates phosphorylation of the CTD (C-terminal domain) of the large subunit of RNA polymerase II (RNAP II) POLR2A (PubMed:[25620562](#), PubMed:[29849146](#)). May play a role in a signaling pathway regulating nuclear functions of cell proliferation (PubMed:[14500717](#)). Modulates alternative splicing by phosphorylating the splice factor SRSF6 (By similarity). Has pro-survival function and negatively regulates the apoptotic process (By similarity). Promotes cell survival upon genotoxic stress through phosphorylation of SIRT1 (By similarity). This in turn inhibits p53/TP53 activity and apoptosis (By similarity). Phosphorylates SEPTIN4, SEPTIN5 and SF3B1 at 'Thr-434' (By similarity).

#### Cellular Location

Nucleus. Nucleus speckle {ECO:0000250|UniProtKB:Q61214}

#### Tissue Location

Ubiquitous. Highest levels in skeletal muscle, testis, fetal lung and fetal kidney.


## Background

DYRK1A is a member of the Dual-specificity tyrosine phosphorylation-regulated kinase (DYRK) family. This member contains a nuclear targeting signal sequence, a protein kinase domain, a leucine zipper motif, and a highly conservative 13-consecutive-histidine repeat. It catalyzes its autophosphorylation on serine/threonine and tyrosine residues. It may play a significant role in a signaling pathway regulating cell proliferation and may be involved in brain development. This gene is a homolog of *Drosophila* *mnb* (minibrain) gene and rat *Dyrk* gene. It is localized in the Down syndrome critical region of chromosome 21, and is considered to be a strong candidate gene for learning defects associated with Down syndrome.

## References

Adayev,T., Biochemistry 46 (25), 7614-7624 (2007)  
 Chang,H.S., Int. J. Cancer 120 (11), 2377-2385 (2007)  
 Alvarez,M., Mol. Biol. Cell 18 (4), 1167-1178 (2007)  
 Wissing,J., Mol. Cell Proteomics 6 (3), 537-547 (2007)

## Images



Anti-DYRK1A Antibody (Center) at 1:2000 dilution + NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 86 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.