

PAK2 Rabbit mAb

Catalog # AP75868

Product Information

Application	WB, IHC-P, FC, IP
Primary Accession	Q13177
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	58043

Additional Information

Gene ID	5062
Other Names	PAK2
Dilution	WB~~1:1000-1:5000 IHC-P~~N/A FC~~1:20 IP~~1:20-1:50
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	PAK2
Function	Serine/threonine protein kinase that plays a role in a variety of different signaling pathways including cytoskeleton regulation, cell motility, cell cycle progression, apoptosis or proliferation (PubMed: 12853446 , PubMed: 16617111 , PubMed: 19273597 , PubMed: 19923322 , PubMed: 33693784 , PubMed: 7744004 , PubMed: 9171063). Acts as a downstream effector of the small GTPases CDC42 and RAC1 (PubMed: 7744004). Activation by the binding of active CDC42 and RAC1 results in a conformational change and a subsequent autophosphorylation on several serine and/or threonine residues (PubMed: 7744004). Full-length PAK2 stimulates cell survival and cell growth (PubMed: 7744004). Phosphorylates MAPK4 and MAPK6 and activates the downstream target MAPKAPK5, a regulator of F-actin polymerization and cell migration (PubMed: 21317288). Phosphorylates JUN and plays an important role in EGF-induced cell proliferation (PubMed: 21177766). Phosphorylates many other substrates including histone H4 to promote assembly of H3.3 and H4 into nucleosomes,

BAD, ribosomal protein S6, or MBP (PubMed:[21724829](#)). Phosphorylates CASP7, thereby preventing its activity (PubMed:[21555521](#), PubMed:[27889207](#)). Additionally, associates with ARHGEF7 and GIT1 to perform kinase-independent functions such as spindle orientation control during mitosis (PubMed:[19273597](#), PubMed:[19923322](#)). On the other hand, apoptotic stimuli such as DNA damage lead to caspase-mediated cleavage of PAK2, generating PAK-2p34, an active p34 fragment that translocates to the nucleus and promotes cellular apoptosis involving the JNK signaling pathway (PubMed:[12853446](#), PubMed:[16617111](#), PubMed:[9171063](#)). Caspase-activated PAK2 phosphorylates MKNK1 and reduces cellular translation (PubMed:[15234964](#)).

Cellular Location

[Serine/threonine-protein kinase PAK 2]: Cytoplasm Nucleus Note=MYO18A mediates the cellular distribution of the PAK2-ARHGEF7-GIT1 complex to the inner surface of the cell membrane


Tissue Location

Ubiquitously expressed. Higher levels seen in skeletal muscle, ovary, thymus and spleen


Background

The activated kinase acts on a variety of targets. Phosphorylates ribosomal protein S6, histone H4 and myelin basic protein. Full length PAK 2 stimulates cell survival and cell growth. The process is, at least in part, mediated by phosphorylation and inhibition of pro-apoptotic BAD.




Images



Western blot analysis of PAK2 in K562, rat Brain, C6, 3T3, Hela lysates using PAK2 antibody.



Immunohistochemistry analysis of paraffin-embedded Human tonsil using PAK2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.