

# Proteasome alpha 6 Rabbit mAb

Catalog # AP75959

## Product Information

---

Application	WB, ICC
Primary Accession	<a href="#">P60900</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	27399

## Additional Information

---

Gene ID	5687
Other Names	PSMA6
Dilution	WB~~1/500-1/1000 ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information



---

Name	PSMA6 ( <a href="#">HGNC:9535</a> )
Synonyms	PROS27
Function	<p>Component of the 20S core proteasome complex involved in the proteolytic degradation of most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP- dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins that could impair cellular functions, and by removing proteins whose functions are no longer required. Associated with the PA200 or PA28, the 20S proteasome mediates ubiquitin- independent protein degradation. This type of proteolysis is required in several pathways including spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented antigenic peptides (20S-PA28 complex).</p>
Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q9QUM9, ECO:0000269 PubMed:12181345}. Nucleus. Note=Translocated from the

cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9 (PubMed:34711951)  
Colocalizes with TRIM5 in cytoplasmic bodies (By similarity)  
{ECO:0000250|UniProtKB:Q9QUM9, ECO:0000269|PubMed:34711951}

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.