

CBX4 Rabbit mAb

Catalog # AP76317

Product Information

Application	WB
Primary Accession	O00257
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	61368

Additional Information

Gene ID	8535
Other Names	CBX4
Dilution	WB~~1:2000-1:5000
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	CBX4
Function	<p>E3 SUMO-protein ligase that catalyzes sumoylation of target proteins by promoting the transfer of SUMO from the E2 enzyme to the substrate (PubMed:12679040, PubMed:22467880, PubMed:22825850). Also acts as a histone reader, which specifically recognizes and binds histone H3 trimethylated at 'Lys-9' and 'Lys-27' (H3K9me3 and H3K27me3, respectively) via its chromo domain (By similarity). Catalyzes sumoylation of HNRNPK, a p53/TP53 transcriptional coactivator, hence indirectly regulates p53/TP53 transcriptional activation resulting in p21/CDKN1A expression (PubMed:22825850). Acts as a regulator of brown adipocyte differentiation by mediating sumoylation of PRDM16, thereby preventing PRDM16 ubiquitination and degradation (By similarity). Monosumoylates ZNF131 (PubMed:22467880). Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:12167701, PubMed:19636380, PubMed:21282530).</p>

PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:[12167701](#), PubMed:[19636380](#), PubMed:[21282530](#)). Plays a role in the lineage differentiation of the germ layers in embryonic development (By similarity).

Cellular Location

Nucleus. Nucleus speckle. Note=Localization to nuclear polycomb bodies is required for ZNF131 sumoylation (PubMed:22467880). Localized in distinct foci on chromatin (PubMed:18927235)


Tissue Location

Ubiquitous.


Background

E3 SUMO-protein ligase which facilitates SUMO1 conjugation by UBE2I. Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility.

Images



Western blot analysis of CBX4 in C6, Raw264.7 lysates using CBX4 antibody.



Western blot analysis of CBX4 in HeLa, A549, HL-60, U251 lysates using CBX4 antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.