

Phospho-Hormone Sensitive Lipase (Ser855) Rabbit mAb

Catalog # AP76344

Product Information

Application WB
Primary Accession Q05469
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 116598

Additional Information

Gene ID 3991

Other Names LIPE

Dilution WB~~1/500-1/1000

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Protein Information

Name LIPE

Function Lipase with broad substrate specificity, catalyzing the hydrolysis of

triacylglycerols (TAGs), diacylglycerols (DAGs), monoacylglycerols (MAGs), cholesteryl esters and retinyl esters (PubMed:15716583, PubMed:15955102, PubMed:19800417, PubMed:8812477). Shows a preferential hydrolysis of DAGs over TAGs and MAGs and preferentially hydrolyzes the fatty acid (FA)

esters at the sn-3 position of the glycerol backbone in DAGs

(PubMed:<u>19800417</u>). Preferentially hydrolyzes FA esters at the sn-1 and sn-2 positions of the glycerol backbone in TAGs (By similarity). Catalyzes the hydrolysis of 2-arachidonoylglycerol, an endocannabinoid and of 2-acetyl monoalkylglycerol ether, the penultimate precursor of the pathway for de novo synthesis of platelet-activating factor (By similarity). In adipose tissue and heart, it primarily hydrolyzes stored triglycerides to free fatty acids, while in steroidogenic tissues, it principally converts cholesteryl esters to free


cholesterol for steroid hormone production (By similarity).

Cellular Location Cell membrane. Membrane, caveola. Cytoplasm, cytosol. Lipid droplet

{ECO:0000250 | UniProtKB:P54310}. Note=Found in the high-density caveolae. Translocates to the cytoplasm from the caveolae upon insulin stimulation (PubMed:17026959). Phosphorylation by AMPK reduces its translocation towards the lipid droplets (By similarity) {ECO:0000250 | UniProtKB:P54310,

ECO:0000269 | PubMed:17026959}

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.