

ERCC8 Rabbit mAb

Catalog # AP76489

Product Information

ApplicationWB, IPPrimary AccessionQ13216ReactivityHumanHostRabbit

Clonality Monoclonal Antibody

Calculated MW 44055

Additional Information

Gene ID 1161

Other Names ERCC8

Dilution WB~~1/500-1/1000 IP~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name ERCC8 {ECO:0000303 | PubMed:19894250, ECO:0000312 | HGNC:HGNC:3439}

Function Substrate-recognition component of the CSA complex, a DCX

(DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complex, involved in transcription-coupled nucleotide excision repair (TC-NER), a process during which RNA polymerase II-blocking lesions are rapidly removed from the transcribed strand of active genes (PubMed:12732143, PubMed:16751180,

PubMed: <u>16964240</u>, PubMed: <u>32142649</u>, PubMed: <u>34526721</u>,

PubMed:38316879, PubMed:38600235, PubMed:38600236). Following recruitment to lesion-stalled RNA polymerase II (Pol II), the CSA complex mediates ubiquitination of Pol II subunit POLR2A/RPB1 at 'Lys- 1268', a critical TC-NER checkpoint, governing RNA Pol II stability and initiating DNA damage excision by TFIIH recruitment (PubMed:12732143, PubMed:16751180,

PubMed:<u>16964240</u>, PubMed:<u>32142649</u>, PubMed:<u>32355176</u>, PubMed:<u>34526721</u>, PubMed:<u>38316879</u>, PubMed:<u>38600235</u>,


PubMed:38600236). The CSA complex also promotes the ubiquitination and subsequent proteasomal degradation of ERCC6/CSB in a UV-dependent manner; ERCC6 degradation is essential for the recovery of RNA synthesis after transcription-coupled repair (PubMed:16751180). Also plays a role in DNA double-strand breaks (DSSBs) repair by non-homologous end joining

(NHEJ) (PubMed: 29545921).

Cellular Location

Nucleus. Chromosome Nucleus matrix. Note=Recruited to lesion- stalled RNA polymerase II (Pol II) sites by ERCC6/CSB (PubMed:32355176). UV-induced translocation to the nuclear matrix is dependent on ERCC6/CSB (PubMed:26620705).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.