

HDAC9 Rabbit mAb

Catalog # AP76522

Product Information

Application WB, IHC-P
Primary Accession Q9UKV0
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 111297

Additional Information

Gene ID 9734

Other Names HDAC9

Dilution WB~~1/500-1/1000 IHC-P~~N/A

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name HDAC9

Synonyms HDAC7, HDAC7B, HDRP, KIAA0744, MITR

Function Responsible for the deacetylation of lysine residues on the N-terminal part

of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses


MEF2-dependent transcription.


Cellular Location Nucleus.

Tissue Location Broadly expressed, with highest levels in brain, heart, muscle and testis.

Isoform 3 is present in human bladder carcinoma cells (at protein level).

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.