

IMPDH2 Rabbit mAb

Catalog # AP76551

Product Information

Application	WB, IHC-P, IHC-F, IP, ICC
Primary Accession	P12268
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	55805




Additional Information

Gene ID	3615
Other Names	IMPDH2
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A IP~~1/20 ICC~~N/A
Format	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	IMPDH2 (HGNC:6053)
Synonyms	IMPD2
Function	Catalyzes the conversion of inosine 5'-phosphate (IMP) to xanthosine 5'-phosphate (XMP), the first committed and rate-limiting step in the de novo synthesis of guanine nucleotides, and therefore plays an important role in the regulation of cell growth (PubMed: 7763314 , PubMed: 7903306). Could also have a single-stranded nucleic acid-binding activity and could play a role in RNA and/or DNA metabolism (PubMed: 14766016). It may also have a role in the development of malignancy and the growth progression of some tumors.
Cellular Location	Cytoplasm. Nucleus. Cytoplasm, cytosol. Note=Can form fiber-like subcellular structures termed 'cytoophidia' in response to intracellular guanine-nucleotide depletion.
Tissue Location	IMPDH1 is the main species in normal leukocytes and IMPDH2 predominates over IMPDH1 in the tumor

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.