

# Ku70 Rabbit mAb

Catalog # AP76564

## Product Information

---

<b>Application</b>	WB, IHC-P, IHC-F, ICC
<b>Primary Accession</b>	<a href="#">P12956</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Calculated MW</b>	69843

## Additional Information

---

<b>Gene ID</b>	2547
<b>Other Names</b>	XRCC6
<b>Dilution</b>	WB~~1/500-1/1000 IHC-P~~N/A IHC-F~~N/A ICC~~N/A
<b>Format</b>	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---

<b>Name</b>	XRCC6 ( <a href="#">HGNC:4055</a> )
<b>Synonyms</b>	G22P1
<b>Function</b>	DNA-binding protein critical for the DNA damage response, specifically in repairing double-strand breaks (DSBs) via the classical non-homologous end joining (NHEJ) pathway. It forms a heterodimer with XRCC5 (Ku80), creating the Ku70:Ku80 heterodimer (Ku complex), which serves as a DNA end-binding complex. It primarily binds DSBs and recruits essential repair factors, assembling the core long-range NHEJ complex to facilitate the alignment and ligation of broken DNA ends (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">33854234</a> , PubMed: <a href="#">34352203</a> , PubMed: <a href="#">9742108</a> ). This pathway ensures the rapid repair of cytotoxic and mutagenic DSBs and contributes to the generation of diversity in T-cell receptors and antibodies through mechanisms such as V(D)J recombination (PubMed: <a href="#">9742108</a> ). Likely acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), catalyzing the beta-elimination of the 5'-deoxyribose- 5-phosphate at abasic sites near DSBs. This activity cleans the termini of abasic sites, a common form of nucleotide damage, preparing broken ends for ligation (PubMed: <a href="#">20383123</a> ). It may also possess 3'-5' DNA helicase activity, although this has not been confirmed in

vivo, and its physiological significance remains unclear (PubMed:[7957065](#)). Beyond DNA repair, the protein contributes to telomere maintenance (PubMed:[29490055](#)). It is also implicated in transcriptional regulation, acting as a cofactor for various transcription factors (PubMed:[12145306](#), PubMed:[8621488](#)). It plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP- RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:[28712728](#)). Can also bind RNAs and recruits PRKDC to a wide range of cellular RNAs, including the U3 small nucleolar RNA, playing a role in the biogenesis of ribosomal RNAs (PubMed:[32103174](#)). Additionally, it negatively regulates apoptosis by interacting with BAX, sequestering it from the mitochondria, and may possess deubiquitination activity targeting BAX (PubMed:[15023334](#), PubMed:[18362350](#), PubMed:[35545041](#)).

### Cellular Location

Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.

### Images

