

MBNL1 Rabbit mAb

Catalog # AP76580

Product Information

ApplicationWBPrimary AccessionQ9NR56ReactivityHumanHostRabbit

Clonality Monoclonal Antibody

Calculated MW 41817

Additional Information

Gene ID 4154

Other Names MBNL1

Dilution WB~~1/500-1/1000

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name MBNL1

Synonyms EXP, KIAA0428, MBNL

Function Mediates pre-mRNA alternative splicing regulation. Acts either as activator

troponin-T (TNNT2) pre-mRNA exon inclusion but induces insulin receptor (IR) pre-mRNA exon inclusion in muscle. Antagonizes the alternative splicing activity pattern of CELF proteins. Regulates the TNNT2 exon 5 skipping through competition with U2AF2. Inhibits the formation of the spliceosome A complex on intron 4 of TNNT2 pre-mRNA. Binds to the stem-loop structure within the polypyrimidine tract of TNNT2 intron 4 during spliceosome assembly. Binds to the 5'-YGCU(U/G)Y-3'consensus sequence. Binds to the IR RNA. Binds to expanded CUG repeat RNA, which folds into a hairpin structure

or repressor of splicing on specific pre-mRNA targets. Inhibits cardiac

containing GC base pairs and bulged, unpaired U residues. Together with RNA binding proteins RBPMS and RBFOX2, activates vascular smooth muscle cells alternative splicing events (PubMed: 37548402). Regulates NCOR2 alternative

splicing (By similarity).


Cellular Location Nucleus. Cytoplasm. Cytoplasmic granule. Note=Localized with DDX1, TIAL1

and YBX1 in stress granules upon stress (PubMed:18335541). Localized in the cytoplasm of multinucleated myotubes (PubMed:18335541). Colocalizes with nuclear foci of retained expanded-repeat transcripts in myotubes from patients affected by myotonic dystrophy (PubMed:10970838, PubMed:11590133, PubMed:11929853)

Tissue Location

Highly expressed in cardiac, skeletal muscle and during myoblast differentiation. Weakly expressed in other tissues (at protein level). Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.