

# NgR3 Rabbit mAb

Catalog # AP76615

#### **Product Information**

**Application** WB. ICC **Q86UN2 Primary Accession** 

Reactivity Human, Mouse, Rat

Host

Clonality Monoclonal Antibody

Calculated MW 49065

#### **Additional Information**

Gene ID 146760

**Other Names** RTN4RL1

**Dilution** WB~~1/500-1/1000 ICC~~N/A

**Format** 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

### **Protein Information**

Name RTN4RL1 ( HGNC:21329)

**Function** Cell surface receptor. Plays a functionally redundant role in postnatal brain

> development and in regulating axon regeneration in the adult central nervous system. Contributes to normal axon migration across the brain midline and normal formation of the corpus callosum. Protects motoneurons against apoptosis; protection against apoptosis is probably mediated by MAG. Plays a role in inhibiting neurite outgrowth and axon regeneration via its binding to neuronal chondroitin sulfate proteoglycans. Binds heparin (By similarity). Like other family members, plays a role in restricting the number dendritic spines and the number of synapses that are formed during brain development (PubMed:<u>22325200</u>). Signaling mediates activation of Rho and downstream

reorganization of the actin cytoskeleton (PubMed:22325200).


**Cellular Location** Cell membrane; Lipid-anchor, GPI-anchor. Membrane raft. Perikaryon


{ECO:0000250 | UniProtKB:Q80WD0}. Cell projection


{ECO:0000250|UniProtKB:Q80WD0}. Note=Localized to the surface of neurons, including axons. {ECO:0000250|UniProtKB:Q80WD0}

**Tissue Location** Predominantly expressed in brain. Expressed at lower levels in kidney, lung,

## **Images**







Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.