

NM23A Rabbit mAb

Catalog # AP76618

Product Information

Application WB
Primary Accession P15531
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 17149

Additional Information

Gene ID 4830

Other Names NME1

Dilution WB~~1/500-1/1000

Format 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

0.05% BSA.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name NME1

Synonyms NDPKA, NM23

Function Major role in the synthesis of nucleoside triphosphates other than ATP. The

ATP gamma phosphate is transferred to the NDP beta phosphate via a ping-pong mechanism, using a phosphorylated active-site intermediate. Possesses nucleoside-diphosphate kinase, serine/threonine-specific protein kinase, geranyl and farnesyl pyrophosphate kinase, histidine protein kinase and 3'-5' exonuclease activities. Involved in cell proliferation, differentiation

and development, signal transduction, G protein-coupled receptor

endocytosis, and gene expression. Required for neural development including neural patterning and cell fate determination. During GZMA- mediated cell death, works in concert with TREX1. NME1 nicks one strand of DNA and TREX1 removes bases from the free 3' end to enhance DNA damage and prevent

DNA end reannealing and rapid repair.

Cellular Location Cytoplasm. Nucleus. Note=Cell-cycle dependent nuclear localization which


can be induced by interaction with Epstein-barr viral proteins or by

degradation of the SET complex by GzmA

Tissue Location

Isoform 1 is expressed in heart, brain, placenta, lung, liver, skeletal muscle, pancreas, spleen and thymus. Expressed in lung carcinoma cell lines but not in normal lung tissues. Isoform 2 is ubiquitously expressed and its expression is also related to tumor differentiation.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.