

# Peroxiredoxin 2 Rabbit mAb

Catalog # AP76651

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, FC                 |
| <b>Primary Accession</b> | <a href="#">P32119</a> |
| <b>Reactivity</b>        | Rat, Human, Mouse      |
| <b>Host</b>              | Rabbit                 |
| <b>Clonality</b>         | Monoclonal Antibody    |
| <b>Isotype</b>           | IgG                    |
| <b>Conjugate</b>         | Unconjugated           |
| <b>Purification</b>      | Affinity Purified      |
| <b>Calculated MW</b>     | 21892                  |

## Additional Information

---

|                    |   |
|--------------------|---|
| <b>Gene ID</b>     | 7001  |
| <b>Other Names</b> | PRDX2   |
| <b>Dilution</b>    | WB~~1:1000 FC~~1:10~50  |
| <b>Format</b>      | Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. |
| <b>Storage</b>     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.        |

## Protein Information

---

|                          |   |
|--------------------------|---|
| <b>Name</b>              | PRDX2   |
| <b>Synonyms</b>          | NKEFB, TDPX1  |
| <b>Function</b>          | Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H <sub>2</sub> O <sub>2</sub> . |
| <b>Cellular Location</b> | Cytoplasm.  |

## Background

---

This gene encodes a member of the peroxiredoxin family of antioxidant enzymes, which reduce hydrogen

peroxide and alkyl hydroperoxides. The encoded protein may play an antioxidant protective role in cells, and may contribute to the antiviral activity of CD8(+) T-cells. This protein may have a proliferative effect and play a role in cancer development or progression.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.