

Superoxide Dismutase 1 Rabbit mAb

Catalog # AP76725

Product Information

Application	WB, IHC-P, FC, IP
Primary Accession	P00441
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Purification	Affinity Purified
Calculated MW	15936

Additional Information

Gene ID	6647
Other Names	SOD1
Dilution	WB~~1:1000 IHC-P~~N/A FC~~1:10~50 IP~~N/A
Format	Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	SOD1 (HGNC:11179)
Function	Destroys radicals which are normally produced within the cells and which are toxic to biological systems (PubMed: 24140062). Catalyzes the oxidation of hydrogen sulfide (H ₂ S) to sulfate, playing an important role in detoxifying H ₂ S and limiting the accumulation of reactive sulfur species (RSS) such as persulfides and polysulfides (PubMed: 36630448).
Cellular Location	Cytoplasm. Nucleus. Note=Predominantly cytoplasmic; the pathogenic variants ALS1 Arg-86 and Ala-94 gradually aggregates and accumulates in mitochondria.

Background

SOD1, Cu/Zn superoxide dismutase, is a major antioxidant enzyme that catalyzes the conversion of

superoxide anion to hydrogen peroxide and molecular oxygen. The mechanism by which mutant SOD1 induces the neurodegeneration observed in ALS is still unclear. Mutant SOD1 proteins become misfolded and consequently oligomerize into high molecular weight species that aggregate and end up in proteinaceous inclusions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.