

# IGF2R Rabbit mAb

Catalog # AP76930

## Product Information

<b>Application</b>	WB, IHC-P, IF, FC, ICC, IP
<b>Primary Accession</b>	<a href="#">P11717</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Immunogen</b>	A synthesized peptide derived from human M6PR
<b>Purification</b>	Affinity Chromatography
<b>Calculated MW</b>	274375

## Additional Information

<b>Gene ID</b>	3482
<b>Other Names</b>	IGF2R
<b>Dilution</b>	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A IP~~N/A
<b>Format</b>	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information



<b>Name</b>	IGF2R
<b>Synonyms</b>	MPRI
<b>Function</b>	Mediates the transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes (PubMed: <a href="#">18817523</a> , PubMed: <a href="#">2963003</a> ). Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex (PubMed: <a href="#">18817523</a> , PubMed: <a href="#">2963003</a> ). The receptor is then recycled back to the Golgi for another round of trafficking through its binding to the retromer (PubMed: <a href="#">18817523</a> ). This receptor also binds IGF2 (PubMed: <a href="#">18046459</a> ). Acts as a positive regulator of T-cell coactivation by binding DPP4 (PubMed: <a href="#">10900005</a> ).

## Cellular Location

Golgi apparatus membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Note=Mainly localized in the Golgi at steady state and not detectable in lysosome (PubMed:18817523) Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface (PubMed:10900005).

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.