

# CHD1L Rabbit mAb

Catalog # AP77109

## Product Information

Application	WB, IHC-P, IF, FC, ICC, IP
Primary Accession	<a href="#">Q86WJ1</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human CHD1L
Purification	Affinity Chromatography
Calculated MW	101000

## Additional Information

Gene ID	9557
Other Names	CHD1L
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

Name	CHD1L {ECO:0000303 PubMed:34210977, ECO:0000312 HGNC:HGNC:1916}
Function	ATP-dependent chromatin remodeler that mediates chromatin- remodeling following DNA damage (PubMed: <a href="#">19661379</a> , PubMed: <a href="#">29220652</a> , PubMed: <a href="#">29220653</a> , PubMed: <a href="#">33357431</a> , PubMed: <a href="#">34210977</a> , PubMed: <a href="#">34486521</a> , PubMed: <a href="#">34874266</a> ). Recruited to DNA damage sites through interaction with poly-ADP-ribose: specifically recognizes and binds histones that are poly-ADP-ribosylated on serine residues in response to DNA damage (PubMed: <a href="#">19661379</a> , PubMed: <a href="#">29220652</a> , PubMed: <a href="#">29220653</a> , PubMed: <a href="#">34486521</a> , PubMed: <a href="#">34874266</a> ). Poly-ADP-ribose-binding activates the ATP-dependent chromatin remodeler activity, thereby regulating chromatin during DNA repair (PubMed: <a href="#">19661379</a> , PubMed: <a href="#">29220652</a> , PubMed: <a href="#">29220653</a> , PubMed: <a href="#">34486521</a> , PubMed: <a href="#">34874266</a> ). Catalyzes nucleosome sliding away from DNA breaks in an ATP-dependent manner (PubMed: <a href="#">19661379</a> , PubMed: <a href="#">29220652</a> , PubMed: <a href="#">29220653</a> ). Chromatin

remodeling activity promotes PARP2 removal from chromatin (PubMed:[33275888](#)).

**Cellular Location**


Nucleus. Chromosome Note=Localizes at sites of DNA damage; recruited by histones H2B and H3 poly-ADP-ribosylated on 'Ser-6' and 'Ser-10', respectively (H2BS6ADPr and H3S10ADPr) by PARP1 or PARP2.

**Tissue Location**

Frequently overexpressed in hepatomacellular carcinomas.

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.