

# Tyrosinase Rabbit mAb

Catalog # AP77225

#### **Product Information**

**Application** WB, IHC-P, IF, ICC

Primary Accession P14679
Reactivity Human
Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human Tyrosinase

**Purification** Affinity Chromatography

Calculated MW 60393

### **Additional Information**

**Gene ID** 7299

Other Names TYR

**Dilution** WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

Name TYR ( HGNC:12442)

**Function** This is a copper-containing oxidase that functions in the formation of

pigments such as melanins and other polyphenolic compounds. Catalyzes the initial and rate limiting step in the cascade of reactions leading to melanin production from tyrosine (By similarity). In addition to hydroxylating tyrosine to DOPA (3,4- dihydroxyphenylalanine), also catalyzes the oxidation of DOPA to DOPA- quinone, and possibly the oxidation of DHI (5,6-dihydroxyindole) to

indole-5,6 quinone (PubMed: 28661582).


**Cellular Location** Melanosome membrane; Single-pass type I membrane protein. Melanosome

{ECO:0000250|UniProtKB:P11344}. Note=Proper trafficking to melanosome is

regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38

{ECO:0000250 | UniProtKB:P11344}

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.