

Cubilin Rabbit mAb

Catalog # AP77394

Product Information

Application WB Primary Accession 060494

Reactivity Rat, Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human Cubilin

Purification Affinity Chromatography

Calculated MW 398736

Additional Information

Gene ID 8029

Other Names CUBN

Dilution WB~~1/500-1/1000

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name CUBN

Synonyms IFCR

Function Endocytic receptor which plays a role in lipoprotein, vitamin and iron

metabolism by facilitating their uptake (PubMed: 10371504, PubMed: 11606717, PubMed: 11717447, PubMed: 14576052,

PubMed:<u>9572993</u>). Acts together with LRP2 to mediate endocytosis of high-density lipoproteins, GC, hemoglobin, ALB, TF and SCGB1A1. Acts together with AMN to mediate endocytosis of the CBLIF-cobalamin complex (PubMed:<u>14576052</u>, PubMed:<u>9572993</u>). Binds to ALB, MB, Kappa and lambdalight chains, TF, hemoglobin, GC, SCGB1A1, APOA1, high density lipoprotein, and the CBLIF-cobalamin complex. Ligand hinding requires calcium.

and the CBLIF-cobalamin complex. Ligand binding requires calcium (PubMed:9572993). Serves as important transporter in several absorptive epithelia, including intestine, renal proximal tubules and embryonic yolk sac. May play an important role in the development of the peri-implantation

embryo through internalization of APOA1 and cholesterol. Binds to LGALS3 at the maternal-fetal interface.


Cellular Location

Apical cell membrane {ECO:0000250|UniProtKB:Q9JLB4}; Peripheral membrane protein. Cell membrane; Peripheral membrane protein {ECO:0000305, ECO:0000305|PubMed:30523278}. Membrane, coated pit. Endosome. Lysosome membrane {ECO:0000250|UniProtKB:O70244}; Peripheral membrane protein. Note=Lacks a transmembrane domain and depends on interaction with AMN for location at the plasma membrane (PubMed:29402915, PubMed:30523278). Colocalizes with AMN and LRP2 in the endocytotic apparatus of epithelial cells (By similarity) {ECO:0000250|UniProtKB:O70244, ECO:0000269|PubMed:29402915, ECO:0000269|PubMed:30523278}

Tissue Location

Detected in kidney cortex (at protein level) (PubMed:9572993). Expressed in kidney proximal tubule cells, placenta, visceral yolk-sac cells and in absorptive intestinal cells. Expressed in the epithelium of intestine and kidney

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.