

CD46 Rabbit mAb

Catalog # AP77413

Product Information

Application WB, IHC-P, IF, ICC

Primary Accession P15529
Reactivity Human
Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human CD46

Purification Affinity Chromatography

Calculated MW 43747

Additional Information

Gene ID 4179

Other Names CD46

Dilution WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name CD46

Synonyms MCP, MIC10

Function Acts as a cofactor for complement factor I, a serine protease which protects

autologous cells against complement-mediated injury by cleaving C3b and

C4b deposited on host tissue. May be involved in the fusion of the

spermatozoa with the oocyte during fertilization. Also acts as a costimulatory factor for T-cells which induces the differentiation of CD4+ into T-regulatory 1

cells. T-regulatory 1 cells suppress immune responses by secreting interleukin-10, and therefore are thought to prevent autoimmunity.

Cellular Location Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane; Single-pass


type I membrane protein. Note=Inner acrosomal membrane of spermatozoa. Internalized upon binding of Measles virus, Herpesvirus 6 or Neisseria gonorrhoeae, which results in an increased susceptibility of infected cells to


complement-mediated injury. In cancer cells or cells infected by Neisseria, shedding leads to a soluble peptide

Tissue Location

Expressed by all cells except erythrocytes.

Images





Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.