

# Rab11A Rabbit mAb

Catalog # AP77443

## Product Information

---

<b>Application</b>	WB, IP
<b>Primary Accession</b>	<a href="#">P62491</a>
<b>Reactivity</b>	Rat, Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal Antibody
<b>Isotype</b>	IgG
<b>Conjugate</b>	Unconjugated
<b>Immunogen</b>	A synthesized peptide derived from human RAB11A
<b>Purification</b>	Affinity Chromatography
<b>Calculated MW</b>	24394

## Additional Information

---

<b>Gene ID</b>	8766
<b>Other Names</b>	RAB11A
<b>Dilution</b>	WB~~1/500-1/1000 IP~~N/A
<b>Format</b>	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information

---


<b>Name</b>	RAB11A ( <a href="#">HGNC:9760</a> )
<b>Function</b>	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed: <a href="#">15601896</a> , PubMed: <a href="#">15689490</a> , PubMed: <a href="#">17462998</a> , PubMed: <a href="#">19542231</a> , PubMed: <a href="#">20026645</a> , PubMed: <a href="#">20890297</a> , PubMed: <a href="#">21282656</a> , PubMed: <a href="#">26032412</a> ). RAB11A regulates endocytic recycling (PubMed: <a href="#">20026645</a> ). Forms a functional Rab11/RAB11FIP3/dynein complex that regulates the movement of peripheral sorting endosomes (SE) along microtubule tracks toward the microtubule organizing center/centrosome, generating the endosomal recycling compartment (ERC) (PubMed: <a href="#">20026645</a> ). Acts as a major regulator of membrane delivery during cytokinesis (PubMed: <a href="#">15601896</a> ). Together with

MYO5B and RAB8A participates in epithelial cell polarization (PubMed:[21282656](#)). Together with Rabin8/RAB3IP, RAB8A, the exocyst complex, PARD3, PRKCI, ANXA2, CDC42 and DNMBP promotes transcytosis of PODXL to the apical membrane initiation sites (AMIS), apical surface formation and lumenogenesis (PubMed:[20890297](#)). Together with MYO5B participates in CFTR trafficking to the plasma membrane and TF (Transferrin) recycling in nonpolarized cells (PubMed:[17462998](#)). Required in a complex with MYO5B and RAB11FIP2 for the transport of NPC1L1 to the plasma membrane (PubMed:[19542231](#)). Participates in the sorting and basolateral transport of CDH1 from the Golgi apparatus to the plasma membrane (PubMed:[15689490](#)). Regulates the recycling of FCGRT (receptor of Fc region of monomeric IgG) to basolateral membranes (By similarity). May also play a role in melanosome transport and release from melanocytes (By similarity). Promotes Rabin8/RAB3IP preciliary vesicular trafficking to mother centriole by forming a ciliary targeting complex containing Rab11, ASAP1, Rabin8/RAB3IP, RAB11FIP3 and ARF4, thereby regulating ciliogenesis initiation (PubMed:[25673879](#), PubMed:[31204173](#)). On the contrary, upon LPAR1 receptor signaling pathway activation, interaction with phosphorylated WDR44 prevents Rab11-RAB3IP-RAB11FIP3 complex formation and cilia growth (PubMed:[31204173](#)). Participates in the export of a subset of neosynthesized proteins through a Rab8-Rab10-Rab11-endosomal dependent export route via interaction with WDR44 (PubMed:[32344433](#)). Also interacts with RABL3 to promote ciliary vesicle formation (PubMed:[36052645](#)).

## Cellular Location

Cell membrane; Lipid-anchor. Endosome membrane; Lipid-anchor. Recycling endosome membrane; Lipid-anchor. Cleavage furrow. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor. Golgi apparatus membrane; Lipid-anchor. Golgi apparatus, trans-Golgi network. Cytoplasmic vesicle membrane; Lipid- anchor. Cell projection, cilium. Note=Localized to WDR44-positive endosomes and tubules (PubMed:[32344433](#)). Translocates with RAB11FIP2 from the vesicles of the endocytic recycling compartment (ERC) to the plasma membrane (PubMed:[11994279](#)). During interphase, localized in vesicles continuously moving from peripheral sorting endosomes towards the pericentrosomal ERC (PubMed:[20026645](#)). Localizes to the cleavage furrow (PubMed:[15601896](#)). Colocalizes with PARD3, PRKCI, EXOC5, OCLN, PODXL and RAB8A in apical membrane initiation sites (AMIS) during the generation of apical surface and lumenogenesis (PubMed:[20890297](#)). Recruited to phagosomes containing *S.aureus* or *M.tuberculosis* (PubMed:[21255211](#)). Localized to rhodopsin transport carriers when interacting with RAB11AFIP3 and ASAP1 in photoreceptors (PubMed:[25673879](#)). Colocalizes with RAB11AFIP1 on punctate vesicles (PubMed:[26032412](#)).

## Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.