

# Tuberin Rabbit mAb

Catalog # AP77484

## Product Information

|                          |  |
|--------------------------|--|
| <b>Application</b>       | WB, IHC-P, FC                                    |
| <b>Primary Accession</b> | <a href="#">P49815</a>                           |
| <b>Reactivity</b>        | Rat, Human, Mouse                                |
| <b>Host</b>              | Rabbit   |
| <b>Clonality</b>         | Monoclonal Antibody                              |
| <b>Isotype</b>           | IgG  |
| <b>Conjugate</b>         | Unconjugated                                     |
| <b>Immunogen</b>         | A synthesized peptide derived from human Tuberin |
| <b>Purification</b>      | Affinity Chromatography                          |
| <b>Calculated MW</b>     | 200608   |

## Additional Information

|                    |  |
|--------------------|--|
| <b>Gene ID</b>     | 7249   |
| <b>Other Names</b> | TSC2   |
| <b>Dilution</b>    | WB~~1/500-1/1000 IHC-P~~N/A FC~~1:10~50  |
| <b>Format</b>      | Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol. |
| <b>Storage</b>     | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.           |

## Protein Information

|                 |  |
|-----------------|--|
| <b>Name</b>     | TSC2 {ECO:0000303   PubMed:7558029, ECO:0000312   HGNC:HGNC:12363}   |
| <b>Function</b> | Catalytic component of the TSC-TBC complex, a multiprotein complex that acts as a negative regulator of the canonical mTORC1 complex, an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed: <a href="#">12172553</a> , PubMed: <a href="#">12271141</a> , PubMed: <a href="#">12842888</a> , PubMed: <a href="#">12906785</a> , PubMed: <a href="#">15340059</a> , PubMed: <a href="#">22819219</a> , PubMed: <a href="#">24529379</a> , PubMed: <a href="#">28215400</a> , PubMed: <a href="#">33436626</a> , PubMed: <a href="#">35772404</a> ). Within the TSC-TBC complex, TSC2 acts as a GTPase- activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1 (PubMed: <a href="#">12172553</a> , PubMed: <a href="#">12820960</a> , PubMed: <a href="#">12842888</a> , PubMed: <a href="#">12906785</a> , PubMed: <a href="#">15340059</a> , PubMed: <a href="#">22819219</a> , PubMed: <a href="#">24529379</a> , PubMed: <a href="#">33436626</a> ). In absence of nutrients, the TSC-TBC complex inhibits mTORC1, thereby preventing phosphorylation of ribosomal protein S6 kinase |

(RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) by the mTORC1 signaling (PubMed:[12172553](#), PubMed:[12271141](#), PubMed:[12842888](#), PubMed:[12906785](#), PubMed:[22819219](#), PubMed:[24529379](#), PubMed:[28215400](#), PubMed:[35772404](#)). The TSC-TBC complex is inactivated in response to nutrients, relieving inhibition of mTORC1 (PubMed:[12172553](#), PubMed:[24529379](#)). Involved in microtubule-mediated protein transport via its ability to regulate mTORC1 signaling (By similarity). Also stimulates the intrinsic GTPase activity of the Ras- related proteins RAP1A and RAB5 (By similarity).

#### Cellular Location



Lysosome membrane; Peripheral membrane protein. Cytoplasm, cytosol  
Note=Recruited to lysosomal membranes in a RHEB-dependent process in absence of nutrients (PubMed:[24529379](#)). In response to insulin signaling and phosphorylation by PKB/AKT1, the complex dissociates from lysosomal membranes and relocates to the cytosol (PubMed:[24529379](#))

#### Tissue Location

Liver, brain, heart, lymphocytes, fibroblasts, biliary epithelium, pancreas, skeletal muscle, kidney, lung and placenta.

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.