

CD10 Rabbit mAb

Catalog # AP77536

Product Information

Application	WB, FC
Primary Accession	P08473
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human CD10
Purification	Affinity Chromatography
Calculated MW	85514


Additional Information

Gene ID	4311
Other Names	MME
Dilution	WB~~1/500-1/1000 FC~~1:10~50
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	MME {ECO:0000303 PubMed:27588448, ECO:0000312 HGNC:HGNC:7154}
Function	Thermolysin-like specificity, but is almost confined on acting on polypeptides of up to 30 amino acids (PubMed: 15283675 , PubMed: 6208535 , PubMed: 6349683 , PubMed: 8168535). Biologically important in the destruction of opioid peptides such as Met- and Leu-enkephalins by cleavage of a Gly-Phe bond (PubMed: 17101991 , PubMed: 6349683). Catalyzes cleavage of bradykinin, substance P and neurotensin peptides (PubMed: 6208535). Able to cleave angiotensin-1, angiotensin-2 and angiotensin 1-9 (PubMed: 15283675 , PubMed: 6349683). Involved in the degradation of atrial natriuretic factor (ANF) and brain natriuretic factor (BNP(1-32)) (PubMed: 16254193 , PubMed: 2531377 , PubMed: 2972276). Displays UV-inducible elastase activity toward skin preelastic and elastic fibers (PubMed: 20876573).
Cellular Location	Cell membrane; Single-pass type II membrane protein

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.