

HDAC8 Rabbit mAb

Catalog # AP77569

Product Information

Application	WB, IHC-P, IF, FC, ICC, IP
Primary Accession	Q9BY41
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human HDAC8
Purification	Affinity Chromatography
Calculated MW	41758

Additional Information

Gene ID	55869
Other Names	HDAC8
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	HDAC8 {ECO:0000303 PubMed:10926844, ECO:0000312 HGNC:HGNC:13315}
Function	Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4) (PubMed: 10748112 , PubMed: 10922473 , PubMed: 10926844 , PubMed: 14701748 , PubMed: 28497810). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed: 10748112 , PubMed: 10922473 , PubMed: 10926844 , PubMed: 14701748). Histone deacetylases act via the formation of large multiprotein complexes (PubMed: 10748112 , PubMed: 10922473 , PubMed: 10926844 , PubMed: 14701748). Also involved in the deacetylation of cohesin complex protein SMC3 regulating release of cohesin complexes from chromatin (PubMed: 22885700). May play a role in smooth muscle cell

contractility (PubMed:[15772115](#)). In addition to protein deacetylase activity, also has protein-lysine deacylase activity: acts as a protein decrotonylase by mediating decrotonylation ((2E)-butenoyl) of histones (PubMed:[28497810](#)).


Cellular Location

Nucleus. Chromosome Cytoplasm Note=Excluded from the nucleoli (PubMed:10748112). Found in the cytoplasm of cells showing smooth muscle differentiation (PubMed:15772115, PubMed:16538051).

Tissue Location

Weakly expressed in most tissues. Expressed at higher level in heart, brain, kidney and pancreas and also in liver, lung, placenta, prostate and kidney.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.