

Factor H Rabbit mAb

Catalog # AP77733

Product Information

Application WB, IF, ICC
Primary Accession P08603
Reactivity Human
Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human Factor H

Purification Affinity Chromatography

Calculated MW 139096

Additional Information

Gene ID 3075

Other Names CFH

Dilution WB~~1/500-1/1000 IF~~1:50~200 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name CFH

Synonyms HF, HF1, HF2

Function Glycoprotein that plays an essential role in maintaining a well-balanced

immune response by modulating complement activation. Acts as a soluble inhibitor of complement, where its binding to self markers such as glycan structures prevents complement activation and amplification on cell surfaces (PubMed:21285368, PubMed:21317894, PubMed:25402769). Accelerates the decay of the complement alternative pathway (AP) C3 convertase C3bBb, thus preventing local formation of more C3b, the central player of the complement

amplification loop (PubMed:19503104, PubMed:21317894,

PubMed: 26700768). As a cofactor of the serine protease factor I, CFH also

regulates proteolytic degradation of already-deposited C3b

(PubMed:<u>18252712</u>, PubMed:<u>23332154</u>, PubMed:<u>28671664</u>). In addition, mediates several cellular responses through interaction with specific

receptors. For example, interacts with CR3/ITGAM receptor and thereby mediates the adhesion of human neutrophils to different pathogens. In turn, these pathogens are phagocytosed and destroyed (PubMed: 20008295, PubMed: 9558116).


Cellular Location

Secreted.

Tissue Location

Expressed in the retinal pigment epithelium (at protein level) (PubMed:25136834). CFH is one of the most abundant complement components in blood where the liver is the major source of CFH protein in vivo. in addition, CFH is secreted by additional cell types including monocytes, fibroblasts, or endothelial cells (PubMed:2139673, PubMed:25136834, PubMed:2968404, PubMed:6444659)

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.