

DNA Polymerase beta Rabbit mAb

Catalog # AP77758

Product Information

Application WB, IHC-P, IF, ICC, IP

Primary Accession <u>P06746</u>

Reactivity Rat, Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human DNA Polymerase beta

Purification Affinity Chromatography

Calculated MW 38178

Additional Information

Gene ID 5423

Other Names POLB

Dilution WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A IP~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name POLB

Function Repair polymerase that plays a key role in base-excision repair

(PubMed: 10556592, PubMed: 9207062, PubMed: 9572863). During this process, the damaged base is excised by specific DNA glycosylases, the DNA backbone is nicked at the abasic site by an apurinic/apyrimidic (AP) endonuclease, and POLB removes 5'-deoxyribose-phosphate from the preincised AP site acting as a 5'-deoxyribose-phosphate lyase (5'-dRP lyase); through its DNA polymerase activity, it adds one nucleotide to the 3' end of the arising single-nucleotide gap (PubMed: 10556592, PubMed: 17526740, PubMed: 9556598, PubMed: 9572863, PubMed: 9614142). Conducts 'gap-filling' DNA synthesis in a stepwise distributive fashion rather than in a processive

fashion as for other DNA polymerases. It is also able to cleave sugar-phosphate bonds 3' to an intact AP site, acting as an AP lyase

(PubMed: 9614142).

Cellular Location

 $\label{lem:nucleus} \textbf{Nucleus. Cytoplasm}. \ \textbf{Note=Cytoplasmic in normal conditions. Translocates to the nucleus following DNA damage}$

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.