

# MVP Rabbit mAb

Catalog # AP77816

### **Product Information**

**Application** WB, IHC-P **Primary Accession** Q14764

**Reactivity** Rat, Human, Mouse

**Host** Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human MVP

**Purification** Affinity Chromatography

Calculated MW 99327

#### **Additional Information**

**Gene ID** 9961

Other Names MVP

**Dilution** WB~~1/500-1/1000 IHC-P~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

**Name** MVP

**Synonyms** LRP

**Function** Required for normal vault structure. Vaults are multi-subunit structures that

may act as scaffolds for proteins involved in signal transduction. Vaults may

also play a role in nucleo-cytoplasmic transport. Down-regulates IFNG-mediated STAT1 signaling and subsequent activation of JAK. Down-regulates SRC activity and signaling through MAP kinases.

**Cellular Location** Cytoplasm. Nucleus, nuclear pore complex. Cytoplasm, perinuclear region.

Note=5% found in the nuclear pore complex (PubMed:15133037). Translocates from the nucleus to the cytoplasm upon EGF treatment


(PubMed:16441665)

**Tissue Location** Present in most normal tissues. Higher expression observed in epithelial cells

with secretory and excretory functions, as well as in cells chronically exposed to xenobiotics, such as bronchial cells and cells lining the intestine.

Overexpressed in many multidrug- resistant cancer cells

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.