

MEIS2 Rabbit mAb

Catalog # AP77957

Product Information

Application WB, FC **Primary Accession** 014770

Reactivity Rat, Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human MEIS2

Purification Affinity Chromatography

Calculated MW 51790

Additional Information

Gene ID 4212

Other Names MEIS2

Dilution WB~~1/500-1/1000 FC~~1:10~50

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name MEIS2

Synonyms MRG1

Function Involved in transcriptional regulation. Binds to HOX or PBX proteins to form

dimers, or to a DNA-bound dimer of PBX and HOX proteins and thought to have a role in stabilization of the homeoprotein-DNA complex. Isoform 3 is required for the activity of a PDX1:PBX1b:MEIS2b complex in pancreatic acinar cells involved in the transcriptional activation of the ELA1 enhancer; the complex binds to the enhancer B element and cooperates with the transcription factor 1 complex (PTF1) bound to the enhancer A element; MEIS2 is not involved in complex DNA-binding. Probably in complex with PBX1, is involved in transcriptional regulation by KLF4. Isoform 3 and isoform 4 can bind to a EPHA8 promoter sequence containing the DNA motif

5'-CGGTCA-3'; in cooperation with a PBX protein (such as PBX2) is proposed to

be involved in the transcriptional activation of EPHA8 in the developing

midbrain. May be involved in regulation of myeloid differentiation. Can bind to the DNA sequence 5'-TGACAG-3'in the activator ACT sequence of the D(1A) dopamine receptor (DRD1) promoter and activate DRD1 transcription; isoform 5 cannot activate DRD1 transcription.

Cellular Location


Nucleus {ECO:0000255 | PROSITE-ProRule:PRU00108}. Cytoplasm, perinuclear

region {ECO:0000250 | UniProtKB:P97367}

Tissue Location

Expressed in various tissues. Expressed at high level in the lymphoid organs of hematopoietic tissues. Also expressed in some regions of the brain, such as the putamen

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.