

# SMARCC1 Rabbit mAb

Catalog # AP77962

## **Product Information**

**Application** WB, IF, FC, ICC, IP

Primary Accession

Reactivity
Rat, Human
Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

Immunogen A synthesized peptide derived from human SMARCC1/BAF155

**Purification** Affinity Chromatography

Calculated MW 122867

# **Additional Information**

**Gene ID** 6599

Other Names SMARCC1

**Dilution** WB~~1/500-1/1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A IP~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

#### **Protein Information**

Name SMARCC1 ( HGNC:11104)

Synonyms BAF155

**Function** Involved in transcriptional activation and repression of select genes by

chromatin remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes that carry out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner. May stimulate the ATPase

activity of the catalytic subunit of the complex (PubMed: 10078207,

PubMed:<u>29374058</u>). Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to

postmitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (By similarity).


**Cellular Location** 

Nucleus. Cytoplasm

**Tissue Location** 

Expressed in brain, heart, muscle, placenta, lung, liver, muscle, kidney and pancreas

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.