

Prostatic Acid Phosphatase Rabbit mAb

Catalog # AP78042

Product Information

Application	WB, IHC-P, IF, ICC, IP
Primary Accession	P15309
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human Prostatic Acid Phosphatase
Purification	Affinity Chromatography
Calculated MW	44566

Additional Information


Gene ID	55
Other Names	ACP3
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	ACP3 (HGNC:125)
Synonyms	ACPP
Function	A non-specific tyrosine phosphatase that dephosphorylates a diverse number of substrates under acidic conditions (pH 4-6) including alkyl, aryl, and acyl orthophosphate monoesters and phosphorylated proteins (PubMed: 10506173 , PubMed: 15280042 , PubMed: 20498373 , PubMed: 9584846). Has lipid phosphatase activity and inactivates lysophosphatidic acid in seminal plasma (PubMed: 10506173 , PubMed: 15280042).
Cellular Location	[Isoform 1]: Secreted
Tissue Location	Highly expressed in the prostate, restricted to glandular and ductal epithelial cells. Also expressed in bladder, kidney, pancreas, lung, cervix, testis and

ovary. Weak expression in a subset of pancreatic islet cells, squamous epithelia, the pilosebaceous unit, colonic neuroendocrine cells and skin adnexal structures. Low expression in prostate carcinoma cells and tissues

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.