

OGA Rabbit mAb

Catalog # AP78081

Product Information

Application	WB, IHC-P, IP
Primary Accession	O60502
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human MGEA5
Purification	Affinity Chromatography
Calculated MW	102915


Additional Information

Gene ID	10724
Other Names	OGA
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	OGA {ECO:0000303 PubMed:20863279, ECO:0000312 HGNC:HGNC:7056}
Function	[Isoform 1]: Cleaves GlcNAc but not GalNAc from O- glycosylated proteins (PubMed: 11148210 , PubMed: 11788610 , PubMed: 20673219 , PubMed: 22365600 , PubMed: 24088714 , PubMed: 28939839 , PubMed: 37962578). Deglycosylates a large and diverse number of proteins, such as CRYAB, ELK1, GSDMD, LMNB1 and TAB1 (PubMed: 28939839 , PubMed: 37962578). Can use p-nitrophenyl-beta-GlcNAc and 4-methylumbelliferone-GlcNAc as substrates but not p-nitrophenyl-beta-GalNAc or p-nitrophenyl-alpha-GlcNAc (in vitro) (PubMed: 20673219). Does not bind acetyl-CoA and does not have histone acetyltransferase activity (PubMed: 24088714).
Cellular Location	[Isoform 3]: Nucleus
Tissue Location	Ubiquitous. Shows highest expression in the brain, placenta and pancreas.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.