

IDH1 Rabbit mAb

Catalog # AP78257

Product Information

Application	WB, IHC-P, IP
Primary Accession	O75874
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human IDH1
Purification	Affinity Purified
Calculated MW	46659


Additional Information

Gene ID	3417
Other Names	IDH1
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	IDH1
Synonyms	PICD
Function	Catalyzes the NADP(+)-dependent oxidative decarboxylation of isocitrate (D-threo-isocitrate) to 2-ketoglutarate (2-oxoglutarate), which is required by other enzymes such as the phytanoyl-CoA dioxygenase (PubMed: 10521434 , PubMed: 19935646). Plays a critical role in the generation of NADPH, an important cofactor in many biosynthesis pathways (PubMed: 10521434). May act as a corneal epithelial crystallin and may be involved in maintaining corneal epithelial transparency (By similarity).
Cellular Location	Cytoplasm, cytosol. Peroxisome

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.