

PAR4 Rabbit mAb

Catalog # AP78315

Product Information

Application WB Primary Accession Q96IZ0

Reactivity Rat, Human, Mouse

Host Rabbit

Clonality Monoclonal Antibody

Isotype IgG

Conjugate Unconjugated

Immunogen A synthesized peptide derived from human PAR4

Purification Affinity Purified

Calculated MW 36568

Additional Information

Gene ID 5074

Other Names PAWR

Dilution WB~~1/500-1/1000

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name PAWR

Synonyms PAR4

Function Pro-apoptotic protein capable of selectively inducing apoptosis in cancer

cells, sensitizing the cells to diverse apoptotic stimuli and causing regression of tumors in animal models. Induces apoptosis in certain cancer cells by activation of the Fas prodeath pathway and coparallel inhibition of

NF-kappa-B transcriptional activity. Inhibits the transcriptional activation and augments the transcriptional repression mediated by WT1. Down-regulates the anti- apoptotic protein BCL2 via its interaction with WT1. Also seems to be a transcriptional repressor by itself. May be directly involved in regulating the

amyloid precursor protein (APP) cleavage activity of BACE1.

Cellular Location Cytoplasm. Nucleus. Note=Mainly cytoplasmic in absence of apoptosis signal


and in normal cells. Nuclear in most cancer cell lines. Nuclear entry seems to

be essential but not sufficient for apoptosis (By similarity). Nuclear localization includes nucleoplasm and PML nuclear bodies.

Tissue Location

Widely expressed. Expression is elevated in various neurodegenerative diseases such as amyotrophic lateral sclerosis, Alzheimer, Parkinson and Huntington diseases and stroke. Down-regulated in several cancers.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.