

Phospho-c-Myc (Thr58) Rabbit mAb

Catalog # AP78552

Product Information

Application	WB, IF, FC, ICC
Primary Accession	P01106
Reactivity	Human
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human Phospho-c-Myc (T58)
Purification	Affinity Chromatography
Calculated MW	50565

Additional Information

Gene ID	4609
Other Names	MYC
Dilution	WB~~1/500-1/1000 IF~~1:50~200 FC~~1:10~50 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information


Name	MYC
Synonyms	BHLHE39
Function	Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed: 24940000 , PubMed: 25956029). Activates the transcription of growth-related genes (PubMed: 24940000 , PubMed: 25956029). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed: 24940000 , PubMed: 25956029). Regulator of somatic reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production

of the PKM M2 isoform (PubMed:[20010808](#)).

Cellular Location

Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm Chromosome.
Note=Association with chromatin is reduced by hyperphosphorylation
(PubMed:30158517) Localization to the nucleolus is dependent on HEATR1
(PubMed:38225354)

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.