

Phospho-Smad5 (Ser463/465) Rabbit mAb

Catalog # AP78566

Product Information

Primary Accession Q99717
Host Rabbit

Clonality Monoclonal Antibody

Calculated MW 52258

Additional Information

Gene ID 4090

Other Names SMAD5

Format Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name SMAD5

Synonyms MADH5

Function Transcriptional regulator that plays a role in various cellular processes

including embryonic development, cell differentiation, angiogenesis and tissue homeostasis (PubMed:12064918, PubMed:16516194). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRIs) and associates with SMAD4 to form a

heteromeric complex which translocates into the nucleus acting as

transcription factor (PubMed:<u>9442019</u>). In turn, the hetero-trimeric complex recognizes cis- regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed:<u>33510867</u>). Non-

phosphorylated SMAD5 has a cytoplasmic role in energy metabolism

regulation by promoting mitochondrial respiration and glycolysis in response to cytoplasmic pH changes (PubMed: 28675158). Mechanistically, interacts

with hexokinase 1/HK1 and thereby accelerates glycolysis


(PubMed: 28675158).

Cellular Location Cytoplasm. Nucleus Mitochondrion. Note=Cytoplasmic in the absence of

ligand. Migrates to the nucleus when complexed with SMAD4

Tissue Location Ubiquitous.

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.