

# FKBP1B Antibody (N-term E6)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7868a

#### **Product Information**

Application WB, IF, E Primary Accession P68106

Other Accession P97534, O8HYX6, O9Z2I2, P68107, P62943, P26883, P62942, P18203

**Reactivity** Human, Mouse **Predicted** Bovine, Rabbit, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB17414
Calculated MW 11783
Antigen Region 1-30

### **Additional Information**

Gene ID 2281

Other Names Peptidyl-prolyl cis-trans isomerase FKBP1B, PPIase FKBP1B, 126 kDa

FK506-binding protein, 126 kDa FKBP, FKBP-126, FK506-binding protein 1B, FKBP-1B, Immunophilin FKBP126, Rotamase, h-FKBP-12, FKBP1B, FKBP126,

FKBP1L, FKBP9, OTK4

Target/Specificity This FKBP1B antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 1-30 amino acids from the N-terminal

region of human FKBP1B.

**Dilution** WB~~1:1000 IF~~1:10~50 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation

followed by dialysis against PBS.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** FKBP1B Antibody (N-term E6) is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Protein Information**

Name FKBP1B ( HGNC:3712)

**Function** Has the potential to contribute to the immunosuppressive and toxic effects

of FK506 and rapamycin. PPIases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in

oligopeptides.

**Cellular Location** Cytoplasm. Sarcoplasmic reticulum

**Tissue Location** Detected in heart muscle (at protein level). Isoform 1 and isoform 2 are

ubiquitous with highest levels in brain and thymus.


# **Background**

FKBP1B is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. This protein is a cis-trans prolyl isomerase that binds the immunosuppressants FK506 and rapamycin. The protein is highly similar to the FK506-binding protein 1A. Its physiological role is thought to be in excitation-contraction coupling in cardiac muscle.


### References

Arakawa H., Nagase H.Biochem. Biophys. Res. Commun. 200:836-843(1994) Lam E., Martin M.M., Timerman A.P.Biol. Chem. 270:26511-26522(1995) Deivanayagam C.C., Carson M.Acta Crystallogr. D 56:266-271(2000)

# **Images**




Western blot analysis of anti-FKBP1B Antibody (N-term E6) (Cat.#AP7868a) in CEM cell line lysates (35ug/lane). FKBP1B (arrow) was detected using the purified Pab.



Western blot analysis of anti-FKBP1B Antibody (N-term E6) (Cat.#AP7868a) in mouse cerebellum tissue lysates (35ug/lane). FKBP1B (arrow) was detected using the purified Pab.

Confocal immunofluorescent analysis of FKBP1B Antibody (N-term E6)(Cat#AP7868a) with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.