NUDT5 Rabbit mAb

Catalog # AP78736

Product Information

Application	WB, IHC-P
Primary Accession	Q9UKK9
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human NUDT5
Purification	Affinity Chromatography
Calculated MW	24328

Additional Information

Gene ID	11164
Other Names	NUDT5
Dilution	WB~~1/500-1/1000 IHC-P~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	NUDT5
Synonyms	NUDIX5 {ECO:0000303|PubMed:27257257}
Function	Enzyme that can either act as an ADP-sugar pyrophosphatase in absence of diphosphate or catalyze the synthesis of ATP in presence of diphosphate (PubMed:27257257). In absence of diphosphate, hydrolyzes with similar activities various modified nucleoside diphosphates such as ADP-ribose, ADP-mannose, ADP-glucose, 8-oxo-GDP and 8-oxo-dGDP (PubMed:10567213, PubMed:10722730, PubMed:17052728, PubMed:19699693, PubMed:21389046). Can also hydrolyze other nucleotide sugars with low activity (PubMed:19699693, PubMed:21389046). In presence of diphosphate, mediates the synthesis of ATP in the nucleus by catalyzing the conversion of ADP-ribose to ATP and ribose 5-phosphate. Nuclear ATP synthesis takes place when dephosphorylated at Thr-45 (PubMed:27257257). Nuclear ATP generation is required for extensive chromatin remodeling events that are

energy-consuming (PubMed:27257257). Does not play a role in U8 snoRNA decapping activity (By similarity). Binds U8 snoRNA (By similarity).

Cellular Location   Nucleus.

Tissue Location   Widely expressed. Most abundant in liver.

Images

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.