

# IL-33 Rabbit mAb

Catalog # AP78822

## Product Information

Application	WB, IHC-P, IF, ICC
Primary Accession	<a href="#">Q8BVZ5</a>
Reactivity	Rat, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human IL-33
Purification	Affinity Chromatography
Calculated MW	29991

## Additional Information

Gene ID	77125
Other Names	IL33
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 ICC~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

## Protein Information


Name	IL33 {ECO:0000312 MGI:MGI:1924375}
Function	Cytokine that binds to and signals through the IL1RL1/ST2 receptor which in turn activates NF-kappa-B and MAPK signaling pathways in target cells (PubMed: <a href="#">29045903</a> ). Involved in the maturation of Th2 cells inducing the secretion of T-helper type 2-associated cytokines (By similarity). Also involved in activation of mast cells, basophils, eosinophils and natural killer cells (By similarity). Acts as an enhancer of polarization of alternatively activated macrophages (By similarity). Acts as a chemoattractant for Th2 cells, and may function as an 'alarmin', that amplifies immune responses during tissue injury (By similarity). Induces rapid UCP2-dependent mitochondrial rewiring that attenuates the generation of reactive oxygen species and preserves the integrity of Krebs cycle required for persistent production of itaconate and subsequent GATA3-dependent differentiation of inflammation-resolving alternatively activated macrophages (PubMed: <a href="#">34644537</a> ).

## Cellular Location

Nucleus. Chromosome {ECO:0000250|UniProtKB:O95760}. Cytoplasm {ECO:0000250|UniProtKB:O95760}. Cytoplasmic vesicle, secretory vesicle {ECO:0000250|UniProtKB:O95760}. Secreted. Note=Secreted and released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore following cleavage by CELA1 (PubMed:35749514, PubMed:35794369). Associates with heterochromatin and mitotic chromosomes (By similarity). The secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10; it results in protein translocation from the cytoplasm into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) followed by vesicle entry and secretion (By similarity). {ECO:0000250|UniProtKB:O95760, ECO:0000269|PubMed:35749514, ECO:0000269|PubMed:35794369}

## Images

---



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.