

CYP1B1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7891c

Product Information

Application	WB, E
Primary Accession	Q16678
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB16944
Calculated MW	60846
Antigen Region	143-176

Additional Information

Gene ID	1545
Other Names	Cytochrome P450 1B1, CYP1B1, CYP1B1
Target/Specificity	This CYP1B1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 143-176 amino acids from the Central region of human CYP1B1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CYP1B1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CYP1B1 {ECO:0000303 PubMed:8910454, ECO:0000312 HGNC:HGNC:2597}
Function	A cytochrome P450 monooxygenase involved in the metabolism of various endogenous substrates, including fatty acids, steroid hormones and vitamins (PubMed: 10681376 , PubMed: 11555828 , PubMed: 12865317 , PubMed: 15258110 , PubMed: 20972997). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second

into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[10681376](#), PubMed:[11555828](#), PubMed:[12865317](#), PubMed:[15258110](#), PubMed:[20972997](#)). Exhibits catalytic activity for the formation of hydroxyestrogens from estrone (E1) and 17beta-estradiol (E2), namely 2- and 4-hydroxy E1 and E2. Displays a predominant hydroxylase activity toward E2 at the C-4 position (PubMed:[11555828](#), PubMed:[12865317](#)). Metabolizes testosterone and progesterone to B or D ring hydroxylated metabolites (PubMed:[10426814](#)). May act as a major enzyme for all-trans retinoic acid biosynthesis in extrahepatic tissues. Catalyzes two successive oxidative transformation of all-trans retinol to all-trans retinal and then to the active form all-trans retinoic acid (PubMed:[10681376](#), PubMed:[15258110](#)). Catalyzes the epoxidation of double bonds of certain PUFA. Converts arachidonic acid toward epoxyeicosatrienoic acid (EpETrE) regioisomers, 8,9-, 11,12-, and 14,15- EpETrE, that function as lipid mediators in the vascular system (PubMed:[20972997](#)). Additionally, displays dehydratase activity toward oxygenated eicosanoids hydroperoxyeicosatetraenoates (HpETEs). This activity is independent of cytochrome P450 reductase, NADPH, and O₂ (PubMed:[21068195](#)). Also involved in the oxidative metabolism of xenobiotics, particularly converting polycyclic aromatic hydrocarbons and heterocyclic aryl amines procarcinogens to DNA-damaging products (PubMed:[10426814](#)). Plays an important role in retinal vascular development. Under hyperoxic O₂ conditions, promotes retinal angiogenesis and capillary morphogenesis, likely by metabolizing the oxygenated products generated during the oxidative stress. Also, contributes to oxidative homeostasis and ultrastructural organization and function of trabecular meshwork tissue through modulation of POSTN expression (By similarity).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q64429}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q64429}. Microsome membrane {ECO:0000250|UniProtKB:Q64429}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q64429}. Mitochondrion {ECO:0000250|UniProtKB:Q64429}. Note=Located primarily in endoplasmic reticulum. Upon treatment with 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), CYP1B1 is also targeted to mitochondria {ECO:0000250|UniProtKB:Q64429}

Tissue Location

Expressed in heart, brain, lung, skeletal muscle, kidney, spleen, thymus, prostate, testis, ovary, small intestine, colon, and peripheral blood leukocytes (PubMed:[8175734](#)). Expressed in retinal endothelial cells and umbilical vein endothelial cells (at protein level) (PubMed:[19005183](#)).


Background

CYP1B1 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. The enzyme localizes to the endoplasmic reticulum and metabolizes procarcinogens such as polycyclic aromatic hydrocarbons and 17beta-estradiol. Mutations in CYP1B1 gene have been associated with primary congenital glaucoma; therefore it is thought that the enzyme also metabolizes a signaling molecule involved in eye development, possibly a steroid.

References

- Suri,F., Mol. Vis. 14, 2349-2356 (2008)
Nelson,D.R., Pharmacogenetics 14 (1), 1-18 (2004)
Stoilov,I., Hum. Mol. Genet. 6 (4), 641-647 (1997)

Images



Western blot analysis of anti-CYP1B1 Antibody (Center)
(Cat.#AP7891c) in CEM cell line lysates (35ug/lane).
CYP1B1(arrow) was detected using the purified Pab.

Citations

- [Evaluation of the effect of the new methoxy-stilbenes on expression of receptors and enzymes involved in estrogen synthesis in cancer breast cells.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.