

# Tyrosine Hydroxylase Rabbit mAb

Catalog # AP78965

#### **Product Information**

**Application** WB, IHC-P, IF, FC, ICC

Primary Accession P07101

**Reactivity** Rat, Human, Mouse

**Host** Rabbit

**Clonality** Monoclonal Antibody

**Isotype** IgG

**Conjugate** Unconjugated

**Immunogen** A synthesized peptide derived from human Tyrosine Hydroxylase

**Purification** Affinity Chromatography

Calculated MW 58600

### **Additional Information**

**Gene ID** 7054

Other Names TH

**Dilution** WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A

Format Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02%

sodium azide and 50% glycerol.

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

### **Protein Information**

Name TH ( <u>HGNC:11782</u>)

**Synonyms** TYH

**Function** Catalyzes the conversion of L-tyrosine to L- dihydroxyphenylalanine

(L-Dopa), the rate-limiting step in the biosynthesis of catecholamines, dopamine, noradrenaline, and adrenaline. Uses tetrahydrobiopterin and molecular oxygen to convert tyrosine to L-Dopa (PubMed:15287903,

PubMed:1680128, PubMed:17391063, PubMed:24753243, PubMed:34922205, PubMed:8528210, Ref.18). In addition to tyrosine, is able to catalyze the hydroxylation of phenylalanine and tryptophan with lower specificity (By similarity). Positively regulates the regression of retinal hyaloid vessels during

postnatal development (By similarity).

Cytoplasm, perinuclear region {ECO:0000250 | UniProtKB:P24529}. Nucleus


{ECO:0000250|UniProtKB:P04177} Cell projection, axon

{ECO:0000250|UniProtKB:P24529}. Cytoplasm {ECO:0000250|UniProtKB:P04177}. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle {ECO:0000250|UniProtKB:P04177}. Note=When phosphorylated at Ser-19 shows a nuclear distribution and when phosphorylated at Ser-31 as well at Ser-40 shows a cytosolic distribution (By similarity). Expressed in dopaminergic axons and axon terminals. {ECO:0000250|UniProtKB:P04177}

**Tissue Location** 

Mainly expressed in the brain and adrenal glands.

## **Images**



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.