

NAT10 Rabbit mAb

Catalog # AP79053

Product Information

Application	WB, IHC-P, IF, FC, ICC, IP
Primary Accession	Q9H0A0
Reactivity	Rat, Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Isotype	IgG
Conjugate	Unconjugated
Immunogen	A synthesized peptide derived from human NAT10
Purification	Affinity Purified
Calculated MW	115730

Additional Information

Gene ID	55226
Other Names	NAT10
Dilution	WB~~1/500-1/1000 IHC-P~~N/A IF~~1:50~200 FC~~1:10~50 ICC~~N/A IP~~N/A
Format	Liquid in 10mM PBS, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information



Name	NAT10 {ECO:0000255 HAMAP-Rule:MF_03211}
Function	RNA cytidine acetyltransferase that catalyzes the formation of N(4)-acetylcytidine (ac4C) modification on mRNAs, 18S rRNA and tRNAs (PubMed: 25411247 , PubMed: 25653167 , PubMed: 30449621 , PubMed: 35679869). Catalyzes ac4C modification of a broad range of mRNAs, enhancing mRNA stability and translation (PubMed: 30449621 , PubMed: 35679869). mRNA ac4C modification is frequently present within wobble cytidine sites and promotes translation efficiency (PubMed: 30449621). Mediates the formation of ac4C at position 1842 in 18S rRNA (PubMed: 25411247). May also catalyze the formation of ac4C at position 1337 in 18S rRNA (By similarity). Required for early nucleolar cleavages of precursor rRNA at sites A0, A1 and A2 during 18S rRNA synthesis (PubMed: 25411247 , PubMed: 25653167). Catalyzes the formation of ac4C in serine and leucine tRNAs (By similarity). Requires the tRNA-binding adapter

protein THUMPD1 for full tRNA acetyltransferase activity but not for 18S rRNA acetylation (PubMed:[25653167](#)). In addition to RNA acetyltransferase activity, also able to acetylate lysine residues of proteins, such as histones, microtubules, p53/TP53 and MDM2, in vitro (PubMed:[14592445](#), PubMed:[17631499](#), PubMed:[19303003](#), PubMed:[26882543](#), PubMed:[27993683](#), PubMed:[30165671](#)). The relevance of the protein lysine acetyltransferase activity is however unsure in vivo (PubMed:[30449621](#)). Activates telomerase activity by stimulating the transcription of TERT, and may also regulate telomerase function by affecting the balance of telomerase subunit assembly, disassembly, and localization (PubMed:[14592445](#), PubMed:[18082603](#)). Involved in the regulation of centrosome duplication by acetylating CENATAC during mitosis, promoting SASS6 proteasome degradation (PubMed:[31722219](#)). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:[34516797](#)).

Cellular Location

Nucleus, nucleolus {ECO:0000255 | HAMAP- Rule:MF_03211, ECO:0000269 | PubMed:12429849, ECO:0000269 | PubMed:14592445, ECO:0000269 | PubMed:19303003, ECO:0000269 | PubMed:24786082, ECO:0000269 | PubMed:25653167, ECO:0000269 | PubMed:30165671, ECO:0000269 | PubMed:34516797}. Midbody {ECO:0000255 | HAMAP-Rule:MF_03211, ECO:0000269 | PubMed:19303003} Note=Nucleolar in interphase and redistributes to the perichromosomal layer and to the midbody during telophase

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.