

# CALM1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP8549b

## Product Information

---

<b>Application</b>	WB, FC, IHC-P, IF, E
<b>Primary Accession</b>	<a href="#">P0DP23</a>
<b>Other Accession</b>	<a href="#">P05419</a> , <a href="#">P62155</a> , <a href="#">P62161</a> , <a href="#">P62160</a> , <a href="#">P62204</a> , <a href="#">P62152</a> , <a href="#">Q6PI52</a> , <a href="#">P62149</a> , <a href="#">O16305</a> , <a href="#">P62157</a> , <a href="#">NP_008819</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Mouse, Rat, Rabbit, Zebrafish, Chicken, Xenopus, Bovine, C.Elegans, Drosophila
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB21103
<b>Calculated MW</b>	16838
<b>Antigen Region</b>	107-132

## Additional Information

---

<b>Gene ID</b>	801;805;808
<b>Other Names</b>	Calmodulin, CaM, CALM1, CALM, CAM, CAM1
<b>Target/Specificity</b>	This CALM1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 107-132 amino acids from the C-terminal region of human CALM1.
<b>Dilution</b>	WB~~1:1000 FC~~1:10~50 IHC-P~~1:100~500 IF~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CALM1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CALM1 {ECO:0000303 PubMed:7925473, ECO:0000312 HGNC:HGNC:1442}
-------------	--

<b>Function</b>	<p>Calmodulin acts as part of a calcium signal transduction pathway by mediating the control of a large number of enzymes, ion channels, aquaporins and other proteins through calcium-binding (PubMed:<a href="#">16760425</a>, PubMed:<a href="#">23893133</a>, PubMed:<a href="#">26969752</a>, PubMed:<a href="#">27165696</a>, PubMed:<a href="#">28890335</a>, PubMed:<a href="#">31454269</a>, PubMed:<a href="#">35568036</a>). Calcium-binding is required for the activation of calmodulin (PubMed:<a href="#">16760425</a>, PubMed:<a href="#">23893133</a>, PubMed:<a href="#">26969752</a>, PubMed:<a href="#">27165696</a>, PubMed:<a href="#">28890335</a>, PubMed:<a href="#">31454269</a>, PubMed:<a href="#">35568036</a>). Among the enzymes to be stimulated by the calmodulin-calcium complex are a number of protein kinases, such as myosin light-chain kinases and calmodulin-dependent protein kinase type II (CaMK2), and phosphatases (PubMed:<a href="#">16760425</a>, PubMed:<a href="#">23893133</a>, PubMed:<a href="#">26969752</a>, PubMed:<a href="#">27165696</a>, PubMed:<a href="#">28890335</a>, PubMed:<a href="#">31454269</a>, PubMed:<a href="#">35568036</a>). Together with CCP110 and centrin, is involved in a genetic pathway that regulates the centrosome cycle and progression through cytokinesis (PubMed:<a href="#">16760425</a>). Is a regulator of voltage- dependent L-type calcium channels (PubMed:<a href="#">31454269</a>). Mediates calcium- dependent inactivation of CACNA1C (PubMed:<a href="#">26969752</a>). Positively regulates calcium-activated potassium channel activity of KCNN2 (PubMed:<a href="#">27165696</a>). Forms a potassium channel complex with KCNQ1 and regulates electrophysiological activity of the channel via calcium- binding (PubMed:<a href="#">25441029</a>). Acts as a sensor to modulate the endoplasmic reticulum contacts with other organelles mediated by VMP1:ATP2A2 (PubMed:<a href="#">28890335</a>).</p>
<b>Cellular Location</b>	<p>Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cell projection, cilium, flagellum {ECO:0000250 UniProtKB:P0DP26}  Note=Distributed throughout the cell during interphase, but during mitosis becomes dramatically localized to the spindle poles and the spindle microtubules</p>

## Background



CALM1 is a member of the EF-hand calcium-binding protein family. Calmodulin mediates the control of a large number of enzymes and other proteins by Ca(2+). Among the enzymes to be stimulated by the calmodulin-Ca(2+) complex are a number of protein kinases and phosphatases. Together with CEP110 and centrin, is involved in a genetic pathway that regulates the centrosome cycle and progression through cytokinesis.

## References


Zhao,D., et.al., Zhonghua Yi Xue Za Zhi 88 (35), 2452-2456 (2008)  
Martins-de-Souza,D., et.al., J. Neural Transm. 116 (3), 275-289 (2009)

## Images


All lanes : Anti-CALM1 Antibody (C-term) at 1:1000 dilution  
Lane 1: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution.  
Observed band size : 17kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.



CALM1 Antibody (C-term) (Cat.#AP8549b) flow cytometry analysis of Hela cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



CALM1 Antibody (C-term) (Cat. #AP8549b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CALM1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of CALM1 Antibody (C-term)(Cat. #AP8549b) with Hela cell followed by Alexa Fluor® 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.