

NLRP3 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8564A

Product Information

Application	WB, IHC-P, E
Primary Accession	Q96P20
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	118173
Antigen Region	65-91

Additional Information

Gene ID	114548
Other Names	NACHT, LRR and PYD domains-containing protein 3, Angiotensin/vasopressin receptor AII/AVP-like, Caterpillar protein 11, CLR11, Cold autoinflammatory syndrome 1 protein, Cryopyrin, PYRIN-containing APAF1-like protein 1, NLRP3, C1orf7, CIAS1, NALP3, PYPAF1
Target/Specificity	This NLRP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 65-91 amino acids from the N-terminal region of human NLRP3.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NLRP3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NLRP3 {ECO:0000303 PubMed:17907925, ECO:0000312 HGNC:HGNC:16400}
Function	Sensor component of the NLRP3 inflammasome, which mediates inflammasome activation in response to defects in membrane integrity,

leading to secretion of inflammatory cytokines IL1B and IL18 and pyroptosis (PubMed:[16407889](#), PubMed:[18403674](#), PubMed:[18604214](#), PubMed:[23582325](#), PubMed:[25686105](#), PubMed:[27929086](#), PubMed:[28656979](#), PubMed:[28847925](#), PubMed:[30487600](#), PubMed:[30612879](#), PubMed:[31086327](#), PubMed:[31086329](#), PubMed:[31189953](#), PubMed:[33231615](#), PubMed:[34133077](#), PubMed:[34341353](#), PubMed:[34512673](#), PubMed:[36442502](#), PubMed:[40450990](#)). In response to pathogens and other damage-associated signals that affect the integrity of membranes, initiates the formation of the inflammasome polymeric complex composed of NLRP3, CASP1 and PYCARD/ASC (PubMed:[16407889](#), PubMed:[18403674](#), PubMed:[27432880](#), PubMed:[28847925](#), PubMed:[31189953](#), PubMed:[33231615](#), PubMed:[34133077](#), PubMed:[34341353](#), PubMed:[36142182](#), PubMed:[36442502](#)). Recruitment of pro-caspase-1 (proCASP1) to the NLRP3 inflammasome promotes caspase-1 (CASP1) activation, which subsequently cleaves and activates inflammatory cytokines IL1B and IL18 and gasdermin-D (GSDMD), promoting cytokine secretion and pyroptosis (PubMed:[23582325](#), PubMed:[28847925](#), PubMed:[31189953](#), PubMed:[33231615](#), PubMed:[34133077](#), PubMed:[34341353](#)). Activation of NLRP3 inflammasome is also required for HMGB1 secretion; stimulating inflammatory responses (PubMed:[22801494](#)). Involved in the homeostatic wound healing response to tissue injury, a multistep cascade that guides neutrophil migration to necrotic sites while avoiding collateral damage of healthy tissues. ATP released from necrotic cells triggers activation of NLRP3 inflammasome through P2RX7 signaling leading to neutrophil adhesion to the vascular endothelium close to the injury site (By similarity). Under resting conditions, ADP-bound NLRP3 is autoinhibited (PubMed:[35114687](#)). NLRP3 activation stimuli include extracellular ATP, nigericin, reactive oxygen species, crystals of monosodium urate or cholesterol, amyloid-beta fibers, environmental or industrial particles and nanoparticles, such as asbestos, silica, aluminum salts, cytosolic dsRNA, etc (PubMed:[16407889](#), PubMed:[18403674](#), PubMed:[18604214](#), PubMed:[19414800](#), PubMed:[23871209](#)). Almost all stimuli trigger intracellular K(+) efflux (By similarity). These stimuli lead to membrane perturbation and activation of NLRP3 (By similarity). Upon activation, NLRP3 is transported to microtubule organizing center (MTOC), where it is unlocked by NEK7, leading to its relocalization to dispersed trans- Golgi network (dTGN) vesicle membranes and formation of an active inflammasome complex (PubMed:[36442502](#), PubMed:[39173637](#)). Associates with dTGN vesicle membranes by binding to phosphatidylinositol 4- phosphate (PtdIns4P) (PubMed:[30487600](#), PubMed:[34554188](#)). Shows ATPase activity (PubMed:[17483456](#)).

Cellular Location

Cytoplasm, cytosol. Inflammasome. Cytoplasm, cytoskeleton, microtubule organizing center. Golgi apparatus membrane. Endoplasmic reticulum {ECO:0000250|UniProtKB:Q8R4B8}. Mitochondrion. Secreted. Nucleus {ECO:0000250|UniProtKB:Q8R4B8} Note=In macrophages, under resting conditions, mainly located in the cytosol and on membranes of various organelles, such as endoplasmic reticulum, mitochondria and Golgi: forms an inactive double-ring cage that is primarily localized on membranes (By similarity). Upon activation, NLRP3 is transported to microtubule organizing center (MTOC), where it is unlocked by NEK7, leading to its relocalization to dispersed trans-Golgi network (dTGN) vesicle membranes for the formation of an active inflammasome complex (PubMed:[39173637](#)) Recruited to dTGN vesicle membranes by binding to phosphatidylinositol 4-phosphate (PtdIns4P) (PubMed:[30487600](#)). After the induction of pyroptosis, inflammasome specks are released into the extracellular space where they can further promote IL1B processing and where they can be engulfed by macrophages (PubMed:[24952504](#)). Phagocytosis induces lysosomal damage and inflammasome activation in the recipient cells (PubMed:[24952504](#)). In the Th2 subset of CD4(+) helper T cells, mainly located in the nucleus (By similarity).

Nuclear localization depends upon KPNA2 (By similarity). In the Th1 subset of CD4(+) helper T cells, mainly cytoplasmic (By similarity).

{ECO:0000250 | UniProtKB:Q8R4B8, ECO:0000269 | PubMed:24952504, ECO:0000269 | PubMed:30487600, ECO:0000269 | PubMed:39173637}

Tissue Location

Predominantly expressed in macrophages (PubMed:33231615, PubMed:34133077). Also expressed in dendritic cells, B- and T-cells (at protein level) (PubMed:11786556, PubMed:17164409) Expressed in LPS-treated granulocytes, but not in resting cells (at protein level) (PubMed:17164409). Expression in monocytes is very weak (at protein level) (PubMed:17164409). Expressed in stratified non-keratinizing squamous epithelium, including oral, esophageal and ectocervical mucosa and in the Hassall's corpuscles in the thymus Also, detected in the stratified epithelium covering the bladder and ureter (transitional mucosa) (at protein level) (PubMed:17164409) Expressed in lung epithelial cells (at protein level) (PubMed:23229815). Expressed in chondrocytes (PubMed:12032915) Expressed at low levels in resting osteoblasts (PubMed:17907925)


Background

NLRP3 is a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leucine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NALP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis.

References

Feldmann, J., et al., Am. J. Hum. Genet. 71 (1), 198-203 (2002)

Images



All lanes : Anti-NLRP3 Antibody (N-term) at 1:500 dilution
Lane 1: A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size : 100kDa Blocking/Dilution buffer: 5% NFDN/TBST.

Citations

- [Exposure to environmental black carbon exacerbates nasal epithelial inflammation via the reactive oxygen species \(ROS\)-nucleotide-binding, oligomerization domain-like receptor family, pyrin domain containing 3 \(NLRP3\)-caspase-1-interleukin 1β \(IL-1β\) pathway](#)
- [Protective effects of catechin against monosodium urate-induced inflammation through the modulation of NLRP3 inflammasome activation.](#)
- [Effects of Berberine on NLRP3 and IL-1β Expressions in Monocytic THP-1 Cells with Monosodium Urate Crystals-Induced Inflammation.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.