

# C5 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP8577a

## Product Information

---

<b>Application</b>	WB, IHC-P, FC, E
<b>Primary Accession</b>	<a href="#">P01031</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB21866
<b>Calculated MW</b>	188305
<b>Antigen Region</b>	253-281

## Additional Information

---

<b>Gene ID</b>	727
<b>Other Names</b>	Complement C5, C3 and PZP-like alpha-2-macroglobulin domain-containing protein 4, Complement C5 beta chain, Complement C5 alpha chain, C5a anaphylatoxin, Complement C5 alpha' chain, C5, CPAMD4
<b>Target/Specificity</b>	This C5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 253-281 amino acids of human C5.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	C5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	C5 {ECO:0000303   PubMed:1984448, ECO:0000312   HGNC:HGNC:1331}
<b>Function</b>	Precursor of the C5a anaphylatoxin and complement C5b components of the complement pathways, which consist in a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens

the adaptive immune system (PubMed:[12878586](#), PubMed:[18204047](#), PubMed:[30643019](#), PubMed:[6554279](#)). Activated downstream of classical, alternative, lectin and GZMK complement pathways (PubMed:[12878586](#), PubMed:[18204047](#), PubMed:[30643019](#), PubMed:[39914456](#), PubMed:[39814882](#), PubMed:[6554279](#)).

## Cellular Location

Secreted. [C5a anaphylatoxin]: Secreted


## Background

C5 is the fifth component of complement, which plays an important role in inflammatory and cell killing processes. This protein is comprised of alpha and beta polypeptide chains that are linked by a disulfide bridge. An activation peptide, C5a, which is an anaphylatoxin that possesses potent spasmogenic and chemotactic activity, is derived from the alpha polypeptide via cleavage with a convertase. The C5b macromolecular cleavage product can form a complex with the C6 complement component, and this complex is the basis for formation of the membrane attack complex, which includes additional complement components.


## References

McClure,A., et.al., *Rheumatology (Oxford)* 48 (11), 1369-1374 (2009)  
DiScipio,R.G.et.al., *J. Biol. Chem.* 267 (24), 17087-17094 (1992)

## Images




Western blot analysis of C5 Antibody (N-term) (Cat. #AP8577a) in HL-60 cell line lysates (35ug/lane). C5 (arrow) was detected using the purified Pab.(2ug/ml)



Formalin-fixed and paraffin-embedded human breast carcinoma with C5 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Flow cytometric analysis of MDA-231 cells using C5 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



## Citations

---

- [SAK-HV Triggered a Short-period Lipid-lowering Biotherapy Based on the Energy Model of Liver Proliferation via a Novel Pathway.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.