

WNT5B Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8603C

Product Information

Application WB, IHC-P, FC, E

Primary Accession Q9H1J7

Other Accession P33945, P22726, P31286, Q90X07, Q27052, P22725, P41221

Reactivity Human, Mouse **Predicted** Rat, Rabbit, Xenopus

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB22511
Calculated MW 40323
Antigen Region 153-182

Additional Information

Gene ID 81029

Other Names Protein Wnt-5b, WNT5B

Target/Specificity This WNT5B antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 153-182 amino acids from the Central

region of human WNT5B.

Dilution WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent

concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions WNT5B Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name WNT5B

Function Ligand for members of the frizzled family of seven transmembrane

receptors. Probable developmental protein. May be a signaling molecule

which affects the development of discrete regions of tissues. Is likely to signal over only few cell diameters (By similarity).

Cellular Location

Secreted, extracellular space, extracellular matrix


Background

WNT5B is a protein which shows 94% and 80% amino acid identity to the mouse Wnt5b protein and the human WNT5A protein, respectively.


References

Saitoh, T. et.al., Int. J. Mol. Med. 10 (3), 345-349 (2002) Yates, K.E. et.al., Osteoarthr. Cartil. 12 (6), 497-505 (2004)


Images



Western blot analysis of WNT5B Antibody (Center) (Cat. #AP8603c) in mouse bladder tissue lysates (35ug/lane). WNT5B (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human skeletal muscle with WNT5B Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of widr cells using WNT5B Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.